

John Barnard
Steven Lund
USPAS
January 19-30, 2015
Hampton, Virginia

Summary of JB lectures

START WITH MICROSCOPIC PHASE SPACE DENSITY

$$N(\underline{x}, \underline{v}, t) = \sum_{i=1}^N \delta(\underline{x} - \underline{x}_i(t)) \delta(\underline{v} - \underline{v}_i(t))$$

Klimontovich Density

$\frac{\partial N}{\partial t} + \text{LIEBMAN EQUATIONS} \Rightarrow$ KLIMONTIVICH EQUATION:

$$\frac{\partial N}{\partial t} + \underline{v} \cdot \nabla_{\underline{x}} N(\underline{x}, \underline{v}, t) - \frac{q}{m} (E^m + \underline{v} \times B^m) \cdot \nabla_{\underline{v}} N(\underline{x}, \underline{v}, t) = 0$$

$$\text{or } \frac{dN(\underline{x}, \underline{v}, t)}{dt} = 0$$

Letting $N = f + \delta f$ $f = \langle N \rangle$
 $E^m = E + \delta E$ $E = \langle E^m \rangle$
 $B^m = B + \delta B$ $B = \langle B^m \rangle$

$$f = \frac{\int N d^3x d^3v}{\Delta x^3 \Delta v^3}$$

$n^{-1/3} \ll \Delta x \ll \lambda_D$

PERFORMING LOCAL AVERAGES TO OBTAIN SMOOTH & "LIKEY" QUANTITIES:

$$\frac{\partial f}{\partial t} + \underline{v} \cdot \frac{\partial f}{\partial \underline{x}} + \frac{d\underline{v}}{dt} \cdot \frac{\partial f}{\partial \underline{v}} = \frac{\partial f}{\partial t_c} \sim \frac{f}{\tau_c}$$

We estimated $\left| \frac{\partial f / \partial t_c}{\left(\frac{qE}{m} \cdot \frac{\partial f}{\partial v} \right)} \right| \sim \frac{1}{16 \lambda_D^3 n_0} \ll 1$

$\lambda_D = v_{th} / \omega_p$ $v_{th} \equiv \sqrt{\frac{kT}{m}}$ $\omega_p \equiv \sqrt{\frac{q^2 n}{\epsilon_0 m}}$

$$\frac{\partial f}{\partial t} + \underline{v} \cdot \frac{\partial f}{\partial \underline{x}} + \frac{d\underline{p}}{dt} \cdot \frac{\partial f}{\partial \underline{p}} = 0 \quad \dot{p} = -\frac{\partial H}{\partial \underline{x}} \quad \dot{x} = \frac{\partial H}{\partial \underline{p}}$$

$$\frac{df}{dt} = 0$$

LIUVILLE'S EQUATION (INCOMPRESSIBILITY OF PHASE VOLUME)

DEFINE NORMALIZED EMITTANCES PROPORTIONAL TO $\frac{\Delta p_x \Delta z}{\Delta x \Delta y} \propto \Delta E \Delta t$

SO THAT $\epsilon_{px}^2 = \gamma_{\beta}^2 (\langle x^2 \rangle \langle x'^2 \rangle - \langle x x' \rangle^2)$

\Rightarrow CONSTANT IF FORCES ARE LINEAR IN x & FILAMENTATION IS ABSENT (LINEAR WITHOUT COUPLING TO z , DC y).

WE DERIVED TWO SETS OF PARTICLE EQUATION OF MOTION:

AXIAL EQUATION (FOR AXISYMMETRIC SYSTEMS) ($\frac{\partial}{\partial \theta} = 0$)

STARTING WITH THE LORENZ FORCE EQUATION $\frac{d\mathbf{p}}{dt} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$ IN CYL. COORD.

$$\frac{d}{dt}(\gamma m r \dot{\theta}) - \gamma m r \dot{\theta}^2 = q \left(\frac{V''}{z} r + r \dot{\theta} B \right) + q (E_r^{\text{self}} + v_z B_{\theta}^{\text{self}})$$

\uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow
 INITIAL CENTRIFUGAL E_r external $v_z B_z$ SELF-FIELDS
 (DIVERGENCE OF $E = 0$)

$\dot{r} \equiv \frac{dr}{dt}; \quad v = \frac{dr}{ds} = \frac{\dot{r}}{\beta c}$

θ -component:

$$p_{\theta} = \gamma m r^2 \dot{\theta} + \frac{q B(z) r^2}{2} = \text{constant}$$

$$= \gamma m r^2 \beta c \theta' + \frac{q B r^2}{2} = \text{constant}$$

$$r'' + \frac{(\gamma \beta)'}{\gamma \beta} r' + \frac{\gamma''}{z \beta^2 \gamma} r + \left(\frac{\omega_c}{z \gamma \beta c} \right)^2 r - \left(\frac{p_{\theta}}{\gamma \beta m c} \right)^2 \frac{1}{r^3} - \frac{q}{\gamma m v_z^2} \frac{\lambda(r)}{2 \pi \epsilon_0 r} = 0$$

\uparrow \uparrow \uparrow \uparrow \uparrow \uparrow \uparrow
 INITIAL ACCELERATION E_r CENTRIFUGAL CENTRIFUGAL SELF-FIELDS
 (INERTIA) (CONVERGENCE OF FIELD LINES)

STATISTICAL AVERAGE OF THIS EQUATION

$$r_b'' + \frac{(\gamma \beta)'}{\gamma \beta} r_b' + \frac{\gamma''}{z \beta^2 \gamma} r_b + \left(\frac{\omega_c}{z \gamma \beta c} \right)^2 r_b - \frac{4 \langle p_{\theta}^2 \rangle}{(\gamma m \beta c)^2 r_b^3} - \frac{E_r}{r_b} - \frac{Q}{r_b} = 0$$

$$E_r^2 \equiv 4(\langle r^2 \rangle \langle r'^2 \rangle - \langle r r' \rangle^2 + \langle r z \rangle \langle r^2 \theta'^2 \rangle - \langle r^2 \theta' \rangle^2); \quad Q = \frac{q \lambda}{2 \pi \epsilon_0 \gamma^3 \beta^2 m c^2}$$

$$= E_x^2 - 4 \langle r^2 \theta'^2 \rangle \quad (\text{if } p = p(r) \text{ only})$$

CARTESIAN EQUATION OF MOTION

J BARWARD

(15)

EQUATION OF MOTION AGAIN STARTING WITH $\frac{d\mathbf{p}}{dt} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$

RETURN TO X, Y COORDINATES

$$x'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) x' = \frac{-q}{\gamma^3 m v_z^2} \frac{\partial \phi}{\partial x} \mp \begin{cases} \frac{q B'}{\gamma m v_z^2} x & \text{An magnetic quads} \\ \frac{q E'}{\gamma m v_z^2} x & \text{An electric quads} \end{cases}$$

Let $\frac{\gamma m v_z}{q} = \frac{p}{q} \equiv [B'] \equiv \text{RIGIDITY}$

$$y'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) y' = \frac{-q}{\gamma^3 m v_z^2} \frac{\partial \phi}{\partial y} \mp \begin{cases} \frac{B'}{[B']} y & \text{magnetic} \\ \frac{q E'}{\gamma m v_z^2} y & \text{electric} \end{cases}$$

Define $r_x, r_y, \epsilon_x, \epsilon_y$ in terms of 2nd order moments

ENVELOPE EQUATION

$$r_x^2 = 4 \langle x^2 \rangle; \quad r_y^2 = 4 \langle y^2 \rangle$$

$$r_x' = \frac{4 \langle x x' \rangle}{r_x}$$

$$r_x'' = \frac{4 \langle x x'' \rangle}{r_x} + \frac{\epsilon_x^2}{r_x^3}; \quad \epsilon_x^2 = 16 (\langle x^2 \rangle \langle x'^2 \rangle - \langle x x' \rangle^2)$$

$$r_y'' = \frac{4 \langle y y'' \rangle}{r_y} + \frac{\epsilon_y^2}{r_y^3}; \quad \epsilon_y^2 = 16 (\langle y^2 \rangle \langle y'^2 \rangle - \langle y y' \rangle^2)$$

for magnetic focusing:

$$r_x'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) r_x' + \frac{4q}{\gamma^3 m v_z^2} \left\langle x \frac{\partial \phi}{\partial x} \right\rangle \mp \frac{B'}{[B']} r_x - \frac{\epsilon_x^2}{r_x^3} = 0$$

$$r_y'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) r_y' + \frac{4q}{\gamma^3 m v_z^2} \left\langle y \frac{\partial \phi}{\partial y} \right\rangle \mp \frac{B'}{[B']} r_y - \frac{\epsilon_y^2}{r_y^3} = 0$$

(for electric focusing $\frac{B'}{[B']} \rightarrow \frac{q E'}{\gamma m v_z^2}$)

SPACE CHARGE TERM WITH ELLIPTICAL SYMMETRY

NOW DEFOCUSING IN ONE DIRECTION AND FOCUSING IN THE OTHER \Rightarrow RADIAL SYMMETRY SHOULD BE REPLACED

BY ELLIPTICAL SYMMETRY:
$$\rho = \rho \left(\frac{x^2}{v_x^2} + \frac{y^2}{v_y^2} \right)$$

CAN BE SHOWN THAT
$$\left\langle x \frac{\partial \phi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{v_x}{v_x + v_y}$$

$$\left\langle y \frac{\partial \phi}{\partial y} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{v_y}{v_x + v_y}$$

USE
$$\phi(x,y) = \frac{-\lambda v_x v_y}{4\epsilon_0} \int_0^{\infty} \frac{\eta(\chi) ds}{\sqrt{v_x^2 + s} \sqrt{v_y^2 + s}}$$
 TO PROVE, WHERE $\hat{\rho}(\chi) = \frac{d\eta}{d\chi}$

$$\rho(x,y) = \hat{\rho}(\chi) \Big|_{\chi=0}$$

$$\chi = \frac{x^2}{v_x^2 + s} + \frac{y^2}{v_y^2 + s}$$

DEFINING
$$Q = \frac{2\lambda q}{4\pi\epsilon_0 \gamma^3 m v_z^2}$$

$$v_x'' + \frac{1}{\gamma v_z^2} \frac{d}{ds} (\gamma v_z) v_x' - \frac{2Q}{v_x + v_y} = \frac{B'}{[B\rho]} v_x - \frac{\epsilon_x^2}{v_x^3} = 0$$

$$v_y'' + \frac{1}{\gamma v_z^2} \frac{d}{ds} (\gamma v_z) v_y' - \frac{2Q}{v_x + v_y} = \frac{B'}{[B\rho]} v_y - \frac{\epsilon_y^2}{v_y^3}$$

(for Electric Focusing $\frac{B'}{[B\rho]} \rightarrow \frac{qE'}{\gamma m v_z^2}$)

(ANALOGUE TO CIRCULAR BEAM:

$$\left\langle r \frac{\partial \phi}{\partial r} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \quad \text{PROVED IN HOMEWORK}$$



Envelope Equations derived in course:

I. Statistical rms envelope equations (envelopes defined in terms of rms quantities; emittance not guaranteed to be a conserved quantity).

1. Paraxial: r_b ; azimuthal symmetry; $\rho(r)$
2. Cartesian; r_x, r_y ; elliptical symmetry $\rho(x^2/r_x^2 + y^2/r_y^2)$
3. Longitudinal: r_z for $E_z = -\frac{g}{4\pi\epsilon_0} \frac{\partial\lambda}{\partial z} \propto z$; $\lambda \propto (1 - 4z^2/r_z^2)$; $v \propto z/r_z$
4. Ellipsoidal (rf) bunches: r_\perp, r_z (Also r_x, r_y, r_x ; cf Wangler sec 9.9)
5. Cartesian with images: r_x, r_y
6. Larmor frame: periodic solenoids: \tilde{r}_x, \tilde{r}_z
7. Cartesian including scattering: r_x, r_y ; emittance evolves $(\frac{d\epsilon_x^2}{ds} = 4C_{sc} r_x^2)$

II. Kinetic envelope equations (constraint equations governing the parameters of the distribution function. Emittance conserved.)

1. KV distribution elliptical uniform density beam
 $f(x, x', y, y') \sim \delta(1 - C_x - C_y)$; $E_x \sim x$; $E_y \sim y$;
(Identical envelope equation to #2 above).
2. Neuffer distribution for 1D (longitudinal projections of phase space) parabolic line charge density profiles
 $f(z, z') \sim (1 - C_z)^{1/2}$; $E_z \sim z$;
(Identical envelope equation to #3 above).

III. Moment equations

1. Transverse with chromatic effects

$$\langle x^2 \rangle, \langle xx' \rangle, \langle x'^2 \rangle, \langle x^2 \delta \rangle, \langle xx' \delta \rangle, \langle x'^2 \delta \rangle, \dots$$

Summary of current limits for different focusing systems

Einzel lens

$$Q_{\max} \approx \frac{3\pi^2}{8} \left(\frac{q\phi_0}{mV_0^2} \right)^2 \left(\frac{r_b}{L} \right)^2$$

Here $2\phi_0$ = voltage between Einzel lenses;

Vq = quad voltage relative to ground; qV = ion energy

For non-relativistic beams: $\lambda_{\max} \approx 4\pi\epsilon_0 VQ_{\max}$

$$\lambda_{\max} \propto \frac{\phi_0^2}{V}$$

$$\lambda_{\max} \propto \frac{q}{m} B^2 r_p^2$$

Solenoid

$$Q_{\max} \approx \left(\frac{\omega_c r_b}{2\gamma\beta c} \right)^2$$

$$Q_{\max} \approx \frac{\eta\sigma_0}{2\pi}$$

Quadrupole

$$\left\{ \begin{array}{l} \left(\frac{\eta\pi}{\sin \frac{\eta\pi}{2}} \right) \left(\frac{Br_b}{[B\rho]} \right) \left(\frac{r_b}{r_p} \right) \text{ magnetic} \\ \left(\frac{\eta\pi}{2} \right) \left(\frac{2qV_q}{\gamma m v_z^2} \right) \left(\frac{r_b}{r_p} \right)^2 \text{ electric} \end{array} \right.$$

$$\sigma_0 \sim \left\{ \begin{array}{l} \eta L^2 B / (r_p [B\rho]) \text{ electric} \\ 2\eta L^2 q V_q / (r_p^2 \gamma m v_z^2) \text{ magnetic} \end{array} \right.$$

$$\lambda_{\max} \propto \left\{ \begin{array}{l} \left(\frac{qV}{m} \right)^{1/2} Br_b \text{ magnetic} \\ V_q \text{ electric} \end{array} \right.$$

For non-relativistic beams: $I_{\max} \approx \beta c \lambda_{\max} = \left(\frac{qV}{m} \right)^{1/2} \lambda_{\max}$


$$I_{\max} \propto \left(\frac{q}{m} \right)^{1/2} \frac{\phi_0^2}{V^{1/2}}$$

$$I_{\max} \propto \left(\frac{q}{m} \right)^{3/2} V^{1/2} B^2 r_p^2$$

$$I_{\max} \propto \left\{ \begin{array}{l} \left(\frac{qV}{m} \right) Br_b \text{ magnetic} \\ \left(\frac{qV}{m} \right)^{1/2} V_q \text{ electric} \end{array} \right.$$



Current Limits from Different Focusing Methods



LONGITUDINAL DYNAMICS Summary

1D VLASOV EQUATION (Vlasov Equation) $dx dx' dy dy'$

$$\frac{\partial \tilde{f}}{\partial t} + z' \frac{\partial f}{\partial z} + z'' \frac{\partial \tilde{f}}{\partial z'} = 0$$

$$z'' = \frac{q E_z}{m v_0^2}$$

$$\frac{\partial^2 \phi}{\partial z^2} + \frac{1}{r} \frac{\partial}{\partial r} (r \frac{\partial \phi}{\partial r}) = - \rho / \epsilon_0$$

$$E_z = -\frac{q}{4\pi\epsilon_0} \frac{\partial \lambda}{\partial z}$$

"g-factor" model

CHILD-LANGMUIR IN 1-D PIPE

LEADS TO FLUID EQUATIONS (1D Vlasov Equation) dz'

$$\frac{\partial \lambda}{\partial t} + \frac{\partial (\lambda z')}{\partial z} = 0$$

$$\frac{\partial z'}{\partial t} + z' \frac{\partial z'}{\partial z} + \frac{1}{\lambda} \frac{\partial (\lambda z' z)}{\partial z} = \frac{q E_z}{m v_0^2}$$

- 1D E_z ⇒ CHILD LANGMUIR SOLUTION ← NON-LINEAR SOLUTION TO FLUID EQUATIONS
- g-factor: ⇒ SPACE-CHARGE WAVES
- ↳ LONGITUDINAL OR PERPENDICULAR WALL INSTABILITY (IF $E_z = z' I_z$)
- ⇒ SPACE-CHARGE CATERFACTION WAVES ← NON-LINEAR SOLUTION TO FLUID EQNS.
- Outward expansion at $2c_s$; Inward at c_s
- ⇒ VARIABIC BUNCH COMPRESSION ← NON-LINEAR SOLUTION TO FLUID EQUATIONS
- ⇒ "EAK" FIELDS

2D PIERCE ELECTRODE
TIME DEPENDENT LAMPEL REFERENCE SOLUTION

VLASOV EQUATION ALSO ⇒ ENVELOPE EQUATION $\int \int$ VLASOV Equation $dz dz'$

$$\frac{d^2 n_z}{dz^2} = \frac{E_z^2}{v_z^3} + \frac{3}{2} \frac{q q Q_c}{4\pi\epsilon_0 m v^2} \frac{1}{v_z^2} - K(r) v_z$$

KINETIC SOLUTION TO VLASOV EQUATION & SATISFYING RWI ENVELOPE EQUATION
i. NEUBER DISTRI BUTION

$$f(z, z') = \frac{3N}{2\pi E_z} \sqrt{1 - \frac{z^2}{v_z^2} - \frac{v_z^2 (z' - v_z^2 / v_z)^2}{E_z^2}}$$

ESTIMATING SLOT SIZE

$$r_x'' + \frac{(Y_0 \beta_0)'}{Y_0 \beta_0} r_x' + K_x r_x - \frac{zQ}{r_x + r_y} - \frac{E_x^2}{v_x^3} = 0$$


$$r_y'' + \frac{(Y_0 \beta_0)'}{Y_0 \beta_0} r_y' + K_y r_y - \frac{zQ}{r_x + r_y} - \frac{E_y^2}{v_y^3} = 0$$

IN CHAMBER: NO EXTERNAL FOCUSING, NO ACCELERATION
AND BEAM IS OFTEN CIRCULAR (BY DESIGN)

$\Rightarrow K_x = K_y = (Y_0 \beta_0)' = 0$ & $v_x = v_y = v_b$

\Rightarrow ENVELOPE EQUATION IS:

$$r_b'' = \frac{Q}{r_b} + \frac{E^2}{v_b^3}$$



MULTIPLYING BY r_b' & INTEGRATING \Rightarrow

$$\frac{r_{bf}^{1/2}}{2} - \frac{r_{b0}^{1/2}}{2} = Q \ln \frac{r_{bf}}{r_{b0}} + \frac{E^2}{2v_{b0}^3} - \frac{E^2}{2v_{bf}^3}$$

Now $r_{b0}' \approx 0$ $r_{bf} = \text{spot radius}$
 $r_{bf}' = 0$ $r_{b0} \approx d \theta$

$r_{bf} \ll r_{b0}$

$$\Rightarrow \theta^2 \approx zQ \ln \left(\frac{d}{r_{bf}} \right) + \frac{E^2}{r_{bf}^2}$$


WHEN $\theta \ll 0$

$$r_{\text{spot}}^2 = \frac{E^2}{\theta^2} + r_{\text{CHROMATIC ABERRATION}}^2 + \dots$$

$$r_{\text{CHROMATIC}}^2 = \alpha^2 d^2 \left(\frac{\theta}{\beta} \right)^2 \theta^2$$

$\alpha \approx 6$ (system dependent)

1. CHROMATIC ABERRATIONS TEND TO BROADEN SPOT




SINCE QUADRUPOLE MAGNET FOCUSING $\propto \frac{1}{v_z}$

(i.e. $x'' = \frac{qB'}{\gamma m v_z} x$) A SPREAD IN LONGITUDINAL VELOCITY GIVES RISE TO A BROADENING OF FINAL SPOT.

$x'' = \frac{qB'}{p} x$ For a single quad:

$$\theta = \Delta x' = \frac{qB'l}{p} x \Rightarrow \frac{d\theta}{dp} = -\frac{qB'l}{p^2} = -\frac{\theta}{p}$$



$$\begin{aligned} r_{spot} &= \theta \delta d \\ &= \theta \frac{dd}{d\theta} \frac{d\theta}{dp} \delta p \\ &= \alpha \theta d \left(\frac{\delta p}{p} \right) \end{aligned}$$

$\alpha =$ some constant depending on focal system

Geometry $\Rightarrow \frac{dd}{d\theta} = \frac{\delta d}{\Delta\theta} = \frac{d}{\theta}$

$$r_{spot} = \theta \frac{dd}{d\theta} \left| \frac{d\theta}{dp} \right| \delta p$$

$$= \theta \left(\frac{d}{\theta} \right) \left(\frac{\theta}{p} \right) \delta p = \theta d \frac{\delta p}{p} \text{ (for a single magnet)}$$

NORMAL MODES

LONGITUDINAL

SPACE-CHARGE WAVES (FLUID)

$$\omega = \pm c_s k \quad [\text{IN COMOVING BEAM FRAME}]$$

$$c_s = \sqrt{\frac{qg\lambda_0}{4\pi\epsilon_0 m}} = \text{SPACE CHARGE WAVE SPEED}$$

TRANSVERSE

ENVELOPE MODES

CONTINUOUS FOCUSING (LONG BUNCHES)

$$\text{BREATHING: } k_B^2 = 2k_{p0}^2 + 2k_p^2$$

$$\text{QUADRUPOLE } k_Q^2 = k_{p0}^2 + 3k_p^2$$

$$\text{(HERE } k_p^2 \equiv k_{p0}^2 - \frac{Q}{F_b^2} \text{)}$$

(ANALOGOUS MODES IN BUNCHED BEAMS)

STUBS LOOKED AT MODES IN PERIODIC SYSTEMS (CONTINUOUS FOCUSING)

+ KINETIC MODES (GLUCKSTEIN MODES)

+ FLUID MODES

INSTABILITIES

1. LONGITUDINAL (RESISTIVE WALL) INSTABILITY

(FLUID INSTABILITY)

2. ELECTRON-ION INSTABILITY

(CENTROID INSTABILITY)

STEVE TALKED ABOUT:

3. ENVELOPE INSTABILITIES

STEVE TALKED ABOUT:

4. KINETIC INSTABILITIES

(DISTRIBUTION FUNCTION DEPENDENT)

5. SINGLE PARTICLE RESONANT INSTABILITIES

- HALO

- RING RESONANCES

Several potential instabilities have been investigated in HIF drivers

Temperature anisotropy instability

After acceleration $T_{\parallel} \ll T_{\perp}$ internal beam modes are unstable; saturation occurs when $\bar{T}_{\parallel} \sim \bar{T}_{\perp}/3$. (cf E.A. Startsev, R.C. Davidson, H. Qin, PRSTAB 6 084401(2003) and references therein).

Longitudinal resistive instability

Module impedance interacts with beam, amplifying space charge waves that are backward propagating in beam frame.

(cf. Reiser, 2nd ed., chap. 6, K. Takayama and R. J. Briggs, eds., in *Induction Accelerators*, [Springer, NY], (2012), chap. 9 and references therein).

Beam-break up (BBU) instability

High frequency waves in induction module cavities interact transversely with beam (cf., K. Takayama and R. J. Briggs, eds., in *Induction Accelerators*, [Springer, NY], (2012), chap. 7 and references therein).

Beam-plasma instability

Beam interacts with residual gas in the target chamber (cf. R.C. Davidson and H. Qin in *Phys. of Intense Charged Particle Beams in High Energy Accelerators*, [Imperial College Press, London], (2001), chap 10).



HALO:

COVE TEST PARTICLE MODEL:

$$x'' = \begin{cases} -\left[k_{p0}^2 - \frac{Q}{v_b^2}\right]x & \text{for } v < v_b \\ -\left[k_{p0}^2 - \frac{Q}{v^2}\right]x & \text{for } v > v_b \end{cases}$$

$$v_b = v_{b0} + \delta v_b \cos(k_B s + \phi)$$

Gluckstein's phase-amplitude analysis:

$$x'' + \overbrace{\left[k_{p0}^2 - \frac{Q}{v_{b0}^2}\right]}^{k_p^2} x = f(x)$$

↑
Non linear + forcing part

$$x = A \sin \psi \quad x' = k_p A \cos \psi \quad \leftarrow \text{PHASE/AMPLITUDE}$$

$\psi = k_p s + \alpha$ If $f=0$ A & α would be constant

$$\Rightarrow A' = \frac{1}{k_p v_{b0}} f \cos \psi \quad \alpha' = -\frac{1}{k_p v_{b0} A} f \sin \psi$$


DEFINE RESONANT PHASE: $\Psi_r = 2\psi - k_B s$


AVERAGE OVER ALL NON-RESONANT FREQUENCIES

$$A_r' = \frac{1}{k_p v_{b0}} \int_{-\pi}^{\pi} f \cos \psi \frac{d\psi}{2\pi}; \quad \alpha_r' = -\frac{1}{k_p A_r} \int_{-\pi}^{\pi} \frac{df}{2\pi} f \sin \psi$$

$\rightarrow A_r', \Psi_r' \rightarrow \omega', \Psi_r' \rightarrow H(\omega', \Psi_r') \rightarrow$ GIVE RESONANT PARTICLE TRAJECTORY & SEPARATRIX

Numerically determined frequency and amplitude of particle oscillations: linear rf focusing





Particle frequency = $(1/2)$ Envelope frequency


As particle dives in and out of envelope, particle is at same phase of envelope oscillation.

Those particles that are exiting the beam when beam radius is small, and entering beam when beam radius is large, get larger "kick" going out and smaller kick coming in, so are driven to large amplitude.


Heavy ion	Residual gas molecule	e^- electron
-----------	-----------------------	----------------

Processes:


1. Coulomb collisions (intra-beam)




2. Coulomb collisions with residual gas




3. Charge exchange




4. Stripping



5. Neutralization




6. Gas Ionization



$$\frac{dn_g}{dt} = \frac{n_g}{\tau} + q_{eff}$$

7. Wall interactions




SUMMARY OF ELECTRON, GAS, PRESSURE, & SCATTERING EFFECTS

- 1. COULOMB COLLISIONS WITHIN BEAM CAN TRANSFER ENERGY FROM I TO II AND PROVIDE LOWER LIMIT ON T_{II} , HIGHER THAN FROM ACCELERATIVE COOLING.
- 2. COULOMB INTERACTIONS WITH RESIDUAL GAS NUCLEI PROVIDE A SOURCE OF EMITTANCE GROWTH (BUT NOT IMPORTANT FOR HIGHER-MAII AND LONG RESIDENCE TIMES).
- 3. PRESSURE INSTABILITY FROM DESOLATION OF RESIDUAL GAS BY STRIKED BEAM IONS HITTING WALL OF BEAM-IONIZED RESIDUAL GAS ATOMS, FORCED TO WALL BY E-FIELD OF BEAM. LIMITS CURRENT IN RINGS OR HIGH-VELOCITY LINAC.
- 4. ELECTRONS CAN CASCADE AND REACH A "QUIET" EQUILIBRIUM (POPULATION OF SIMILAR LINE CHARGE TO THE ION BEAM). ELECTRON-ION TWO STREAM INSTABILITY IS UNSTABLE, AND CAN LEAD TO TRANSVERSE INSTABILITY, SIMILAR TO WHAT IS OBSERVED IN SOME PROTON RINGS.


42-182 100 SHEETS
National Brand
Made in U.S.A.

Summary of

Induction acceleration for HIF consists of several subsystems and a variety of beam manipulations



Inertial fusion energy (IFE) power plants of the future will consist of four parts



A power plant driver would fire about five targets per second to produce as much electricity as today's 1000 Megawatt power plant



The Heavy Ion Fusion Virtual National Laboratory

