

PROBLEM SET 6

LUND/BARNARD
2015 USPAS

Problem ①

Let the equation of motion for a test particle oscillating in a mismatched beam be modeled by:

$$x'' + k_p^2 x = -\frac{Q}{v_{b0}^6} x^5 + \frac{2eQ}{v_{b0}^2} x \cos k_B s$$

Let $\frac{x}{v_{b0}} = A \sin \psi$ & $\frac{x'}{v_{b0}} = k_p A \cos \psi$

where $\psi = k_p s + \alpha$

a) CALCULATE A' & α' IN TERMS OF $A, \psi,$ & $k_B s$, (and parameters).

Let $\Psi_r' = 2k_p - k_B + 2\alpha_r'$

b). AVERAGE OVER RAPID VARIATIONS TO OBTAIN EQUATIONS FOR THE AMPLITUDE AND PHASE OF THE NEAR-RESONANT PARTICLES. (i.e. find $A_r'(\Psi_r, A_r)$ & $\Psi_r'(\Psi_r, A_r)$).

c). DEFINE $w = A_r^2$. FIND $w'(\Psi_r, w)$ & $\Psi_r'(\Psi_r, w)$.

d). FIND THE HAMILTONIAN H , such that

$$w' = \frac{\partial H(w, \Psi_r)}{\partial \Psi_r} \quad \text{and} \quad \Psi_r' = -\frac{\partial H(w, \Psi_r)}{\partial w}$$

e). Verify that $H(w, \Psi_r)$ is a constant.

HINT FOR PART (b): $\sin^6 x = \frac{1}{32} (10 - 15 \cos 2x + 6 \cos 4x - \cos 6x)$
 $\cos x \sin^5 x = \frac{1}{32} (5 \sin 2x - 4 \sin 4x + \sin 6x)$

PROBLEM 2

1. SUPPOSE the space charge electric field exterior to the beam went as

$$E_r \sim \frac{\lambda}{4\pi\epsilon_0} \frac{v_{b0}^2}{r^3} \quad \text{instead of} \quad E_r \sim \frac{\lambda}{4\pi\epsilon_0 r}.$$

Would you expect the radial extent of a beam halo due to mismatch to be larger or smaller than the one obeying the $1/r$ dependence?

HINT: CONSIDER THE LOCATION OF THE RADIUS WHERE THE PARTICLE - ENVELOPE RADIUS IS SMALLEST. (ASSUME HALO EXTENT IS PROPORTIONAL TO THIS RADIUS).

Problem 3

TCE Problem 1

S.M. Lund P1/

Envelope Radii

Envelope Radius of a Nonuniform Density Beam

For a uniform density elliptical beam with envelope radii r_x and r_y

$$n(x,y) = \begin{cases} \frac{\lambda}{2\pi r_x r_y} & ; \quad \frac{x^2}{r_x^2} + \frac{y^2}{r_y^2} \leq 1 \\ 0 & ; \quad \text{otherwise} \end{cases}$$

$\lambda = \text{line-charge density} = \text{const.}$

and we showed in previous problems that:

$$r_x = Z \langle x^2 \rangle_L^{1/2}$$


$$r_y = Z \langle y^2 \rangle_L^{1/2}$$

where $\langle x^2 \rangle = \frac{\int d^2x x^2 n(x,y)}{\int d^2x n(x,y)}$

If the density profile is elliptical with

$$n(x,y) = n(\zeta) \quad ; \quad \zeta^2 \equiv \frac{x^2}{r_{xe}^2} + \frac{y^2}{r_{ye}^2} \quad \begin{matrix} r_{xe} = \text{const} \\ r_{ye} = \text{const} \end{matrix}$$

such that $n(\zeta)$ is monotonic decreasing in ζ with a sharp cutoff at $\zeta = 1$



show that

$$r_{xe} > r_x \equiv Z \langle x^2 \rangle_L^{1/2}$$


$$r_{ye} > r_y \equiv Z \langle y^2 \rangle_L^{1/2}$$

where $\langle x^2 \rangle_L$ and $\langle y^2 \rangle_L$ are defined from the nonuniform density beam. You may use steps/transforms from previous problems.

B/ "Edge" Radius Factor in a 1D Beam

For a 1D sheet beam with uniform density,

$$n(x) = \begin{cases} \frac{N}{2x_b}, & -x_b \leq x \leq x_b \\ 0, & \text{otherwise} \end{cases} \quad N = \text{const}$$



Find an edge radius coefficient F such that

$$x_b = F \langle x^2 \rangle^{1/2}$$

where in 1D phase-space

$$\langle \dots \rangle = \frac{\int_{-\infty}^{\infty} dx' \int_{-\infty}^{\infty} dx \dots f}{\int_{-\infty}^{\infty} dx' \int_{-\infty}^{\infty} dx f}$$

$$f = f(x', x; s)$$

1D distribution

Compare F to the corresponding results for a centered 2D elliptical beam

$$r_x = 2 \langle x^2 \rangle^{1/2}$$

$$r_y = 2 \langle y^2 \rangle^{1/2}$$

Should you expect $F=2$? Why?


Problem 4

TCE Problem 2

S.M. Lund PZ/

2/ Image Charges on a Cylindrical Pipe


Consider a perfectly conducting pipe of radius r_p :



A/ Show that the formula for a line-charge λ at the origin in free-space is:

$$\phi(r) = \frac{-\lambda}{2\pi\epsilon_0} \ln r + \text{const}$$

B/ Use the formula in part A/ to show that a solution to the interior problem $|\vec{x}_1| < r_p$ can be found for a line charge λ at coordinate $x = x_0$ by superimposing the direct charge and an image charge at $x = Cx_0$. Calculate C for cylindrical geometry.



Images can be superimposed to obtain the Green's Function for the 2D calculation of ϕ within the cylinder.

Problem 5,
TCE Problem 8

S.M. Lund PV

Free Expansion of a Beam Envelope

In the absence of applied focusing forces:

$$\Gamma_x'' - \frac{2Q}{\Gamma_x + \Gamma_y} - \frac{\epsilon_x^2}{\Gamma_x^3} = 0$$

Initial conditions
($s=s_i$)

$$\Gamma_y'' - \frac{2Q}{\Gamma_x + \Gamma_y} - \frac{\epsilon_y^2}{\Gamma_y^3} = 0$$

$$\Gamma_x(s=s_i) = \Gamma_{xi}$$

$$\Gamma_y(s=s_i) = \Gamma_{yi}$$

$$\Gamma_x'(s=s_i) = \Gamma_{xi}'$$

$$\Gamma_y'(s=s_i) = \Gamma_{yi}'$$

Part I: $Q=0$, $\epsilon_x \neq 0$, free expansion
without space charge.

a) Show that the envelope Hamiltonian satisfies:

$$\frac{\Gamma_x''}{2} + \frac{\epsilon_x^2}{2\Gamma_x^2} = \text{const.}$$

Γ_y eqn analysis
is same.

b) Show that the equation in a) can be written as an integral from the initial condition to solve for the free expansion as a function of s . You do not need to carry out the integration explicitly.

Part II $\epsilon_x = 0$, $Q \neq 0$ free expansion
without emittance

c) Show using $\Gamma_{\pm} \equiv (\Gamma_x \pm \Gamma_y)/2$
that the coupled envelope equations reduce to:
 $\Gamma_+'' - \frac{Q}{\Gamma_+} = 0$, $\Gamma_-'' = 0$

and satisfy

$$\frac{\Gamma_+''}{2} - Q \ln \Gamma_+ = \text{const}$$

$$\Gamma_- = C_1 s + C_2$$

$C_{1,2}$ constants

TCE Problem 8

P19/

d) Argue that for finite $Q \neq 0$ that the free expansion without emittance will be more rapid than the free expansion without space charge ($Q=0, E_x, E_y \neq 0$) when the beam expands sufficiently.