

# PROBLEM SET 8


## PROBLEM

1. A mass 200 ion beam has an injection energy  $qV = 1 \text{ MeV}$ , a pulse duration =  $10 \text{ ns}$ , a normalized transverse emittance of  $1 \text{ mm-mrad}$ , and a fractional longitudinal momentum spread  $\frac{\Delta p}{p} = 10^{-3}$ . ( $\beta_0 = \sqrt{\frac{2qV}{m_0 c^2}} \approx 0.0033$ ).

Assume the transverse and longitudinal normalized emittance is conserved, and assume that in the final focus region the beam is neutralized, with a spot size determined by the emittance and chromatic effects only. (TAKE THE LONGITUDINAL NORMALIZED EMITTANCE TO BE  $\Delta p / p$ , WHERE  $l_b = \text{length of bunch}$ .)

$$r_{\text{spot}}^2 \approx \frac{\epsilon^2}{\theta^2} + \alpha^2 d^2 \theta^2 \left( \frac{\Delta p}{p} \right)^2 \quad \text{Let } \alpha = 6$$

Here  $\epsilon$  = the unnormalized emittance,  $d$  is the distance between the end of the last magnet and the focal spot, and  $\theta$  is the half angle of the convergent beam.



a) What is the optimum focusing angle  $\theta$  which minimizes the spot radius, (expressed in terms of  $\epsilon$ ,  $d$ , &  $\Delta p/p$ )?

What is the radius of the spot

IF the final ion energy were:  
(Assume  $d = 6 \text{ m}$  and final pulse duration =  $10 \text{ ns}$ ).

b)  $10 \text{ GeV}$ ?

c)  $1 \text{ GeV}$ ?

d) UNDER THE ASSUMPTIONS OF THIS PROBLEM, SHOW THAT  $r_{\text{spot}} \sim 1/\beta^n$  where  $n$  is a positive real number, and find  $n$ .  
(Non-relativistic dynamics may be assumed)

## 1/ Moment Equations and Conservation Constraints

The nonrelativistic Vlasov equation is:

$$\left\{ \frac{\partial}{\partial t} + \vec{v} \cdot \frac{\partial}{\partial \vec{x}} + \frac{q}{m} [\vec{E} + \vec{v} \times \vec{B}] \cdot \frac{\partial}{\partial \vec{v}} \right\} f(\vec{x}, \vec{v}, t) = 0$$

Define a fluid density  $n$  and a fluid flow velocity  $\vec{V}$  by

$$n(\vec{x}, t) = \int d^3v \cdot f(\vec{x}, \vec{v}, t)$$

$$n(\vec{x}, t) \vec{V}(\vec{x}, t) = \int d^3v \vec{v} f(\vec{x}, \vec{v}, t)$$

a) Operate on the Vlasov equation with

$$\int d^3v \dots$$

to derive the continuity equation:

$$\frac{\partial}{\partial t} n(\vec{x}, t) + \frac{\partial}{\partial \vec{x}} \cdot (n(\vec{x}, t) \vec{V}(\vec{x}, t)) = 0$$

b) Can the continuity equation be solved by itself if you specify the initial density field  $n(\vec{x}, t=0)$ ? Why?

c) Operate on Vlasov's equation with

$$\int d^3v \vec{v} \dots$$

to derive the fluid force equation.

SKIP  
DO NOT  
WORK

## TKS/ST Problem 1

S.M. Lund <sup>Plg/</sup>

$$\frac{\partial}{\partial t}(n\vec{V}) + \nabla \cdot (n \langle \vec{v}\vec{v} \rangle) = \frac{q}{m} n (\vec{E} + \vec{V} \times \vec{B})$$

$$\langle \vec{v}\vec{v} \rangle = \int d^3v \vec{v}\vec{v} f / \int d^3v f$$

John Barnard in earlier lectures made a definition of a pressure tensor as

$$\underline{P} = m \int d^3v (\vec{v} - \vec{V})(\vec{v} - \vec{V}) f(\vec{x}, \vec{v}, t) \\ = mn \langle \vec{v}\vec{v} \rangle - mn \vec{V}\vec{V}$$

In terms of this the fluid force eqn can be expressed as:

$$\frac{\partial}{\partial t} \vec{V} + \vec{V} \cdot \frac{\partial}{\partial \vec{x}} \vec{V} = \frac{q}{m} (\vec{E} + \vec{V} \times \vec{B}) - \frac{1}{mn} \frac{\partial}{\partial \vec{x}} \cdot \underline{P}$$

This form is often used in fluid/plasma analysis.

- d) If the continuity and force equation derived in parts a) and c) are analyzed, can they be solved in principle if you specify the initial density field  $n(\vec{x}, t=0)$  and the velocity field  $\vec{V}(\vec{x}, t=0)$ ? Why? Does the answer change if we assume a cold initial beam with  $\underline{P} = 0$ ? Why?

- e) Let  $G(f)$  be some smooth, differentiable function of  $f$  satisfying  $G(f \rightarrow 0) = 0$ . Show that

$$\int d^3x \int d^3v G(f) = \text{const.}$$

with  $G$  specified

This so-called "generalized entropy" measure<sup>a</sup> can be used to check Vlasov simulations. For example:

$$G(f) = f: \int d^3x \int d^3v f = \text{const} \Rightarrow \text{charge cons.}$$

$$G(f) = f^2: \int d^3x \int d^3v f^2 = \text{const} \Rightarrow \text{"enstrophy" cons.}$$

⋮

# TKS Problem 2 Problem 3,

S.M. Lund P 21

## Gluckstern Modes on a kV Beam

2/  $n=1$  Gluckstern mode and the kV envelope equation for the breathing mode.  
 $r_b =$  equilibrium matched beam radius.

a) The Gluckstern mode eigenfunction is given by

$$\delta\phi_n = \begin{cases} \frac{A_n}{2} \left[ P_{n-1}\left(1 - \frac{2r^2}{r_b^2}\right) + P_n\left(1 - \frac{2r^2}{r_b^2}\right) \right] & ; 0 \leq r < r_b \\ 0 & ; r_b < r \leq r_p \end{cases}$$

$$n = 1, 2, 3, \dots ; P_n(x) = \text{nth order Legendre Polynomial}$$

Write down the eigenfunction as an explicit polynomial in  $r$  for  $n=1$  and plot this solution.

### Legendre Polynomials

$$P_0(x) = 1$$

$$P_1(x) = x$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1)$$

b) Apply the Poisson equation:

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial \delta\phi_n}{\partial r} \right) = \frac{-q}{\epsilon_0} \delta n_n(r)$$

to calculate the perturbed mode density  $\delta n_n$  for  $\delta\phi_n$  as a function of  $r$  for  $0 \leq r < r_b$ . (the "body-wave" component). Plot this result.

c) Use part b) to calculate the amount of charge introduced into the system by the "body-wave" perturbation  $\delta n_n(r)$  for  $0 \leq r < r_b$ . How far would the beam edge radius  $r_e = r_b + \delta r_b$  need to change to conserve charge to linear order in  $A_1$ ?

## TKS Problem 2

S.M. Lund p29

- d) Obtain the  $n=1$  Gluckstern mode dispersion relation from the general  $n$  formula presented in class:

$$Z_n + \frac{1 - (\delta/\delta_0)^2}{(\delta/\delta_0)^2} \left[ B_{n+1} \left( \frac{j\omega/kv_0}{\delta/\delta_0} \right) - B_n \left( \frac{j\omega/kv_0}{\delta/\delta_0} \right) \right] = 0$$

From the definitions in the class notes for the  $B_n$  we have:

$$B_0(\omega) = 1$$

$$B_1(\omega) = \frac{(\omega/kv_0)^2}{(\omega/kv_0)^2 - 1}$$

Solve for the mode eigenfrequency  $\omega$  as a function of  $k_{\parallel 0}$  and  $\delta/\delta_0$ .

$k_{\parallel}$  is a spatial wavenumber that we sometimes call a "frequency"

- e) Compare the wavenumber  $k_{\parallel}$  calculated in part d) with the "breathing" envelope mode on a round KV equilibrium where we showed that the mode wavenumber is

$$k_{\parallel \text{ envelope}} = \sqrt{2k_{\perp 0}^2 + 2k_{\perp 0}^2 (\delta/\delta_0)^2}$$

Are the wavenumbers the same? Is it reasonable to identify these as the same modes? (Explain why.)  
Would you expect that the lowest order modes of a kinetic theory to always reproduce the KV envelope modes to lowest order? (Explain why.)