

Mathematics of the Falling Cat

Rajan Mehta

Pennsylvania State University

February 2, 2012













- In these pictures, it appears that the cat is rotating its body.
- The laws of physics say that angular momentum must be conserved.
- Cats can't violate the laws of physics.



Question

How can a cat flip its body without angular momentum?

- Rademaker, Ter Braak (1935) - first solution
- Kane, Scher (1969) - more realistic class of solutions
- Montgomery (1993) - full mathematical theory

The mathematical cat


A cat's body is modeled as a pair of equal cylinders, connected by a joint (its spine). The spine can bend, but it does not twist.




The cat's shape

The **shape** of the cat is given by two angles (ψ, θ) .

- ψ is the angle between the two halves of the cat's body.
- θ describes the direction of the cat's legs ($\theta = 0$ when the front and back legs are closest to each other). A change in θ corresponds to a rotation of the cat's body around the "spinal axis".





1 is $(\psi, \theta) = (\pi/2, 0)$.

2 is $(\psi, \theta) = (3\pi/2, \pi)$.

3 might be $(\psi, \theta) = (2\pi/3, \pi/4)$.

What about 4?

1




2



3




4




Cat dynamics

How does the cat move?

- No angular momentum: If the cat doesn't change its shape, then it will not rotate.
- If the cat changes its shape, then the entire body will rotate to “cancel out” the angular momentum of the shape change.
- We can consider changes in ψ and θ separately.



- A change in ψ is “balanced”: the front and back halves of the body have opposite angular momentum.
- The cat can change ψ without causing the body to rotate.



- As θ changes, the front and back halves of the body are both rotating about the bent spine.
- The total angular momentum vector is parallel to the y -axis.
- The *size* of the total angular momentum depends on ψ .
- The rate of rotation needed to compensate is

$$\frac{\alpha \sin(\psi/2)}{\cos^2(\psi/2) + \alpha \sin^2(\psi/2)}$$


How the cat does it



- 1 It bends forward.
- 2 It swings its legs around until they are positioned correctly (note that its back is arched at this point).
- 3 It is now free to curve its back and prepare for landing.



The Kane-Scher solution



Question

Can you think of a way to drop a cat so it can't land on its feet?

Thanks. (And thanks to Eric Kuehne for the cat drawings)