

PHY422/820: Classical Mechanics

FS 2020

Homework #6 (Due: Oct 16)


October 9, 2020

Problem H11 – Double Pendulum

[15 Points] Consider a planar double pendulum with lengths l_1, l_2 and masses m_1, m_2 (see figure).

1. Construct the Lagrangian of the system and derive the equations of motion.
2. What are the normal frequencies for small angles ($\phi_1, \phi_2 \ll 1$)?

HINT: Use the ansatz $\phi_k = \phi_{0,k} \exp(i\omega t)$ with identical ω for $k = 1, 2$, and determine ω such that the resulting system of equations has a non-trivial solution for $\phi_{0,1}, \phi_{0,2}$! Consider the energy conservation for oscillators to analyze whether terms containing $\dot{\phi}_i$ can be omitted or not.



Problem H12 – Geometry of Central-Force Trajectories

In general, an observed trajectory $\vec{r}(s)$ can result from a multitude of underlying forces. For central forces, however, we can derive a differential equation that determines the force law from $\vec{r}(s)$. Let us explore this in the following.

1. Use the equations of motion for a mass m in a central force field to prove the following differential equation:

$$f(r) = \frac{l^2}{mr^4} \left[\frac{d^2r}{d\phi^2} - \frac{2}{r} \left(\frac{dr}{d\phi} \right)^2 - r \right], \quad (1)$$

where

$$f(r) = -\frac{\partial}{\partial r} V(r). \quad (2)$$

2. Use Eq. (1) to show that trajectories of the form

$$r(\phi) = \frac{p}{1 + \epsilon \cos \phi} \quad (3)$$

with positive constants p and ϵ are generated by a central force of the form

$$f(r) = -\frac{\kappa}{r^2}, \quad \kappa > 0. \quad (4)$$

How is κ related to the other constants in the problem? Sketch or plot the trajectories for $\epsilon = 0$, $0 < \epsilon < 1$, $\epsilon = 1$, and $\epsilon > 1$.

3. Determine the force field that yields trajectories of the form

$$r(\phi) = \frac{r_0 \sqrt{1 - \epsilon^2}}{\sqrt{1 + \epsilon \cos 2\phi}}. \quad (5)$$

Sketch or plot the trajectory for appropriate values for $0 < \epsilon < 1$. How is this trajectory distinct from (3), assuming that the meaning of ϵ is the same?

4. Compute the central force that makes the mass m move on the spiral trajectory

$$r(\phi) = r_0 \exp(-\phi). \quad (6)$$