

PHY422/820: Classical Mechanics

FS 2020

Homework #9 (Due: Nov 6)

October 31, 2020

Problem H18 – The Group of Rotations $SO(3)$

[10 points] Rotations in \mathbb{R}^3 can be represented by **special** 3×3 **orthogonal matrices**, which have the following properties

$$\det \mathbf{R} = 1, \quad \mathbf{R}\mathbf{R}^T = \mathbf{R}^T\mathbf{R} = \mathbf{1}, \quad \mathbf{R}^T = \mathbf{R}^{-1}. \quad (1)$$

Show that these matrices form a group by proving that the following axioms are satisfied:

1. The product $\mathbf{R}_3 = \mathbf{R}_1\mathbf{R}_2$ of two rotation matrices $\mathbf{R}_1, \mathbf{R}_2 \in SO(3)$ is also a rotation matrix, $\mathbf{R}_3 \in SO(3)$.
2. There exists a **neutral element** $\mathbf{E} \in SO(3)$ such that $\mathbf{E}\mathbf{R} = \mathbf{R}\mathbf{E} = \mathbf{R}$ for all $\mathbf{R} \in SO(3)$.
3. For each $\mathbf{R} \in SO(3)$ there exists an **inverse element** $\mathbf{R}^{-1} \in SO(3)$ which satisfies $\mathbf{R}^{-1}\mathbf{R} = \mathbf{R}\mathbf{R}^{-1} = \mathbf{E}$.

Now we relax the condition on the determinant and consider the more general group of orthogonal matrices $O(3)$.

4. Show that the determinant of a real orthogonal matrix can only be $\det \mathbf{O} = \pm 1$.
5. Show that the orthogonal matrices with $\det \mathbf{O} = -1$ do *not* form a group.

HINT:

$$\det \mathbf{AB} = \det \mathbf{A} \det \mathbf{B}, \quad \det \mathbf{A}^{-1} = \frac{1}{\det \mathbf{A}}, \quad \det \mathbf{A}^T = \det \mathbf{A}$$

Problem H19 – Infinitesimal Rotations and SO(3) Generators

[10 points] A counter-clockwise rotation by an angle ϕ around the axis \vec{n} can be expressed in vector form as

$$\vec{r}' = \vec{r} \cos \phi + \vec{n}(\vec{n} \cdot \vec{r})(1 - \cos \phi) + (\vec{n} \times \vec{r}) \sin \phi. \quad (2)$$

1. Show that for infinitesimal angles

$$\vec{r}' = \vec{r} + (\epsilon \vec{n}) \times \vec{r} = (\mathbb{1} + \epsilon) \vec{r}, \quad (3)$$

where we have defined

$$\epsilon \equiv \Phi(\epsilon \vec{n}). \quad (4)$$

2. Use the mapping between vectors and antisymmetric matrices to show that

$$\epsilon = \epsilon n_x \mathbf{L}_x + \epsilon n_y \mathbf{L}_y + \epsilon n_z \mathbf{L}_z, \quad (5)$$

where

$$\mathbf{L}_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \mathbf{L}_y = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad \mathbf{L}_z = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (6)$$

are the so-called **generators** of infinitesimal rotations.

3. Show that the generators satisfy

$$[\mathbf{L}_x, \mathbf{L}_y] = \mathbf{L}_z, \quad [\mathbf{L}_y, \mathbf{L}_z] = \mathbf{L}_x, \quad [\mathbf{L}_z, \mathbf{L}_x] = \mathbf{L}_y, \quad (7)$$

where the commutator is defined as

$$[\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}. \quad (8)$$

4. The generators can be used to construct arbitrary antisymmetric matrices. Show that the matrix exponential of any antisymmetric matrix is a rotation matrix, i.e.,

$$(e^{\mathbf{A}})^T = (e^{\mathbf{A}})^{-1}, \quad \det e^{\mathbf{A}} = 1. \quad (9)$$

The matrix exponential is defined by the series

$$e^{\mathbf{A}} = \sum_{k=0}^{\infty} \frac{1}{k!} \mathbf{A}^k. \quad (10)$$

Problem H20 – Tensors

[10 points] A **rank- n tensor** on \mathbb{R}^3 is an n -tuple of real numbers which has the following behavior under rotations:

$$T'_{i_1 \dots i_n} = \sum_{j_1, \dots, j_n=1}^3 R_{i_1 j_1} \cdots R_{i_n j_n} T_{j_1 \dots j_n}, \quad i_k, j_k = 1, \dots, 3, \quad (11)$$

where $R \in SO(3)$.

1. Show that the scalar product of two arbitrary vectors $\vec{a}, \vec{b} \in \mathbb{R}^3$ is invariant under rotations is a scalar, i.e., a rank-0 tensor. Interpret this result geometrically.
2. Show that the moment of inertia tensor

$$I_{ij} = \int d^3r \rho(\vec{r}) (\vec{r}^2 \delta_{ij} - r_i r_j) \quad (12)$$

is a rank-2 tensor in the sense of Eq. (11).

HINT: To prove that d^3r is a scalar, consider how the volume elements is related to the unit vectors.