

John Barnard
Steven Lund
USPAS
January 19-30, 2015
Hampton, Virginia

Current limits

- A. Axisymmetric
 - 1. Solenoids
 - 3. Einzel lens

- B. Quadrupolar
 - 1. Derivation of envelope equations with elliptic symmetry
 - 2. Current limit using fourier transform method
 - 3. Alternative methods

Yesterday we derived the "Paraxial Ray Equation:"

$$r'' + \frac{(\gamma\beta)'}{\gamma\beta} r' + \frac{\gamma''}{2\gamma\beta^2} r + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r + \left(\frac{p_\theta}{\gamma\beta mc}\right)^2 \frac{1}{r^3} - \frac{q}{\gamma^3 m \beta^2 c^2} \frac{\lambda(r)}{2\pi\epsilon_0 r} = 0$$

↑
Inertial
↑
 E_r from
converging
field lines
↑
↑
Part of
centrifugal
term
↑
Self-field
($E_r^{self} - v_z B_\theta^{self}$)

Accelerative
damping (of
angle r')
Solenoidal
focusing
($v_\theta B_z -$ part
of centrifugal
term)

which together with the conservation of canonical angular momentum,

$$p_\theta \equiv \gamma\beta mc r^2 \theta' + \frac{m\omega_c r^2}{2}$$

and initial conditions, specify the orbit of a particle in an axisymmetric field.


Taking statistical moments, we derived the radial envelope equation.

$$r_b'' + \frac{(\gamma\beta)'}{\gamma\beta} r_b' + \frac{\gamma''}{2\gamma\beta^2} r_b + \left(\frac{\omega_c}{2\gamma\beta c}\right)^2 r_b - \left(\frac{2\langle p_\theta \rangle}{\gamma\beta mc}\right)^2 \frac{1}{r_b^3} - \frac{\epsilon_r^2}{r_b^3} - \frac{Q}{r_b} = 0$$

where

$$\epsilon_r^2 = 4 \left(\langle r^2 \rangle \langle r'^2 \rangle - \langle r r' \rangle^2 + \langle r^2 \rangle \langle r^2 \theta'^2 \rangle - \langle r^2 \theta' \rangle^2 \right)$$

Periodic Einzel lenses



PERIODIC SOLENOIDS





Figure 3.2. Schematic of magnet sets producing a periodic focusing solenoidal field with axial periodicity length S . In Fig. 3.2 (a), successive coils are spaced by S and have the same current polarity $+I_1, +I_1, \dots$. In Fig. 3.2 (b), successive coils are spaced by $S/2$ and have alternating current polarities $+I_1, -I_1, +I_1, \dots$.

(FIGURE FROM
DAVIDSON & QIN,
2003) P. 55
"PHYSICS OF
INTENSE CHARGED
PARTICLE BEAMS
IN HIGH ENERGY
ACCELERATORS"

SOLENOIDAL FOCUSING



Let $\gamma' = \gamma'' = 0$


FOR MAXIMUM TRANSPORT $P_0 = 0$ & $E_r^z = 0$

$$\Rightarrow r_b'' + \left(\frac{\omega_c}{2\gamma\beta c} \right)^2 r_b = \frac{Q}{r_b}$$

FOR A MATCHED BEAM:

$$Q_{max} = \left(\frac{\omega_c}{2\gamma\beta c} \right)^2 r_b^2$$

HEURISTICALLY:



$v_0 = \omega r$

$$\omega^2 r + Q m v^2 \left(\frac{r}{r_b^2} \right) = \underbrace{q \omega r^2 B}_{\uparrow \text{MAGNETIC FORCE INWARD}}$$

↑
centrifugal force

↑
SPACE CHARGE FORCE

↑
MAGNETIC FORCE INWARD

$$\Rightarrow \omega^2 + \frac{Q V^2}{r_b^2} = \omega \omega_c$$

$\omega \omega_c - \omega^2 = \text{MAXIMUM WHEN } \omega = \frac{\omega_c}{2}$

$$\Rightarrow Q_{max} = \left(\frac{\omega_c^2}{4} \right) \left(\frac{r_b^2}{V^2} \right)$$

100% RECYCLED FIBER
100% RECYCLED PAPER
100% RECYCLED WHITE
INNOVATION
© 2010 HALLS OF RECORD


International Brand

SOLENOIDAL FOCUSING - CONTINUED

IN REALITY BEAM ACQUIRES v_{θ} AS BEAM ENTERS SOLENOID:

CAMPAD

CONSIDER SIMPLE STEP FUNCTION APPROXIMATION TO SOLENOID FIELD:



LET $B_z = B_0 \left[\Theta(z) + \Theta(l_m - z) - 1 \right]$ = $\begin{cases} 0 & z < 0 \\ B_0 & 0 < z < l_m \\ 0 & z > l_m \end{cases}$

$\frac{\partial B_z}{\partial z} = B_0 \left[\delta(z) + \delta(l_m - z) \right]$

MEM $\Theta(z) = \begin{cases} 1 & z > 0 \\ 0 & z < 0 \end{cases}$


As we found earlier $\nabla \cdot B = 0 \Rightarrow$



$B_r(r, z) \simeq -\frac{r}{z} \frac{\partial B_z}{\partial z} + \dots = -\frac{r}{z} B_0 \left[\delta(z) + \delta(l_m - z) \right]$

$\Delta p_{\theta}^* = q \int v_z B_r dt = \int_{-\infty}^{0+\epsilon} q B_r dz = -\frac{ngB_0}{z}$

$\Rightarrow v_{\theta} = r \frac{qB_0}{zm} = \frac{rv_c}{z}$

Schematic of Einzel lens





$$V \Rightarrow v_z, \quad V'' \Rightarrow E_r \Rightarrow \Delta r$$

EINZEL LENS - ANALYSIS (DERIVATION FROM ED LEE)

NOW, LET $w_0 = \langle p_0 \rangle = E_r^2 = 0$

$$\Rightarrow r_b'' + \frac{\gamma'}{\beta^2 \gamma} r_b' + \frac{\gamma''}{2\beta^2 \gamma} r_b - \frac{Q}{r_b} = 0$$

ALSO ASSUME $\beta \ll 1$, NON-RELATIVISTIC BEAM $\begin{cases} \gamma' \approx \beta \beta' \\ \gamma'' \approx \beta'^2 + \beta'' \beta \end{cases}$

$$r_b'' + \frac{\beta'}{\beta} r_b' + \left[\frac{1}{2} \frac{\beta'^2}{\beta^2} + \frac{1}{2} \frac{\beta''}{\beta} \right] r_b - \frac{Q}{r_b} = 0$$

To eliminate r_b' term try substitution

$$r_b = \left(\frac{\beta_0}{\beta} \right)^{1/2} R$$

$$r_b' = \left(\frac{\beta_0}{\beta} \right)^{1/2} R' - \frac{1}{2} \left(\frac{\beta}{\beta_0} \right)^{-3/2} R \beta'$$

$$r_b'' = \left(\frac{\beta_0}{\beta} \right)^{1/2} R'' - \left(\frac{\beta}{\beta_0} \right)^{3/2} \frac{R'}{\beta_0} \beta' + \frac{3}{4} \left(\frac{\beta}{\beta_0} \right)^{5/2} \frac{R}{\beta_0^2} \beta'^2 - \frac{1}{2} \left(\frac{\beta}{\beta_0} \right)^{3/2} \frac{R}{\beta_0} \beta''$$

$$\Rightarrow \left(\frac{\beta_0}{\beta} \right)^{1/2} R'' + \frac{3}{4} \left(\frac{\beta}{\beta_0} \right)^{5/2} \frac{\beta'^2}{\beta_0^2} R = \frac{Q}{R} \left(\frac{\beta}{\beta_0} \right)^{1/2}$$

$$\Rightarrow \boxed{R'' = \frac{Q}{R} \left(\frac{\beta}{\beta_0} \right) - \frac{3}{4} \left(\frac{\beta'}{\beta} \right)^2 R}$$

EINZEL LENS - CONTINUUM

MODEL: LET $\phi = \phi_0 \cos\left(\frac{\pi z}{L}\right)$

$$\frac{1}{2} m v^2 + q \phi = \text{constant}$$

$$\Rightarrow v^2 = v_0^2 - \frac{2q\phi}{m} \cos\left(\frac{\pi z}{L}\right)$$

$$v' = \frac{q\phi_0}{m v} \left(\frac{\pi}{L}\right) \sin\left(\frac{\pi z}{L}\right)$$

IF $\left(\frac{2q\phi_0}{m}\right) < c v_0^2$: $\left(\frac{\beta'}{\beta}\right)^2 \approx \left(\frac{q\phi_0}{m v_0}\right)^2 \left(\frac{\pi}{L}\right)^2 \sin^2\left(\frac{\pi z}{L}\right)$

$$R'' = \frac{Q}{R} \left(\frac{\beta}{\beta_0}\right) - \frac{3}{4} \left(\frac{\beta'}{\beta}\right)^2 R$$

FOR EQUILIBRIUM LOOK AT D.C. COMPONENT: $\sin^2(kz) = \frac{1}{2} - \frac{1}{2} \cos 2x$

$$R'' = 0 \Rightarrow \frac{Q}{R} = \frac{3}{4} \left(\frac{\beta'}{\beta}\right)^2 R$$

$$R \bar{E} \left(\frac{\beta}{\beta_0}\right)^{1/2} r_b \Rightarrow \bar{R} = r_b$$

$$\left(\frac{\beta'}{\beta}\right)^2 = \frac{1}{2} \left(\frac{q\phi_0}{m v_0}\right)^2 \left(\frac{\pi}{L}\right)^2$$

$$\Rightarrow Q_{\max} = \frac{3\pi^2}{8} \left(\frac{q\phi_0}{m v_0}\right)^2 \left(\frac{r_b}{L}\right)^2$$


J. BANWAL
 (7/5)

BEAM OPTICS AND FOCUSING SYSTEMS WITHOUT SPACE CHARGE

FROM REISER, p. 112

$$E_x = -E'x$$

$$E_y = E'y$$



$$F_x = -qE'x$$

$$F_y = qE'y$$

ELECTROSTATIC QUADS


Figure 3.15. Electrodes and force lines in an electrostatic quadrupole.

$$B_x = B'y$$

$$B_y = B'x$$

$$F_x = -qV_z B'x$$

$$F_y = qV_z B'y$$



MAGNETIC QUADS

ENVELOPE EQUATIONS FOR NON-AXISYMMETRIC SYSTEMS

(25)

$$r_x^2 \equiv 4 \langle x^2 \rangle \quad r_y^2 \equiv 4 \langle y^2 \rangle$$

$$2 r_x r_x' = 8 \langle x x' \rangle$$

$$r_x' = \frac{4 \langle x x' \rangle}{r_x}$$

$$\begin{aligned} r_x'' &= \frac{4 \langle x x'' \rangle}{r_x} + \frac{4 \langle x'^2 \rangle}{r_x} - \frac{4 \langle x x' \rangle}{r_x^2} r_x' \\ &= \frac{4 \langle x x'' \rangle}{r_x} + \frac{16 \langle x'^2 \rangle \langle x^0 \rangle}{r_x^2} - \frac{16 \langle x x' \rangle^2}{r_x^2} \end{aligned}$$

DEFINE $E_x^2 = 16 (\langle x'^2 \rangle \langle x^0 \rangle - \langle x x' \rangle^2)$

$$\Rightarrow \boxed{r_x'' = \frac{4 \langle x x'' \rangle}{r_x} + \frac{E_x^2}{r_x^3}}$$

SO HOW DO WE CALCULATE $\langle x x'' \rangle$?

RETURN TO SINGLE PARTICLE EQUATION (IN CARTESIAN COORDINATES)

$$\frac{d}{dt} (\gamma m \dot{x}) = \gamma m \ddot{x} = q (E_x + \dot{y} B_z - \dot{z} B_y)$$

\downarrow
 x''
+ similarly
 y''

\downarrow
QUADRUPOLE FOCUSING
SPACE-CHARGE OF ELLIPTICAL
BEAMS

EQUATION OF MOTION

RETURN TO X, Y COORDINATES

$$x'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) x' = \frac{-q}{\gamma^3 m v_z^2} \frac{\partial \phi}{\partial x} \mp \begin{cases} \frac{q B'}{\gamma m v_z^2} x & \text{for magnetic quadrupoles} \\ \frac{q E'}{\gamma m v_z^2} x & \text{for electric quadrupoles} \end{cases}$$

Let $\frac{\gamma m v_z}{q} = \frac{P}{q} \equiv [B'] \equiv \text{RIGIDITY}$

$$y'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) y' = \frac{-q}{\gamma^3 m v_z^2} \frac{\partial \phi}{\partial y} \mp \begin{cases} \frac{B'}{[B']} y & \text{magnetic} \\ \frac{q E'}{\gamma m v_z^2} y & \text{electric} \end{cases}$$

ENVELOPE EQUATION

$$r_x^2 = 4 \langle x^2 \rangle; \quad r_y^2 = 4 \langle y^2 \rangle$$

$$r_x' = \frac{4 \langle x x' \rangle}{r_x}$$

$$r_x'' = \frac{4 \langle x x'' \rangle}{r_x} + \frac{E_x^2}{v_x^3};$$

$$E_x^2 = 16 (\langle x^2 \rangle \langle x'^2 \rangle - \langle x x' \rangle^2)$$

$$r_y'' = \frac{4 \langle y y'' \rangle}{r_y} + \frac{E_y^2}{v_y^3}$$

$$E_y^2 = 16 (\langle y^2 \rangle \langle y'^2 \rangle - \langle y y' \rangle^2)$$

for magnetic focusing:

$$r_x'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) r_x' + \frac{4q}{\gamma^3 m v_z^2} \left\langle x \frac{\partial \phi}{\partial x} \right\rangle \mp \frac{B'}{[B']} r_x - \frac{E_x^2}{v_x^3} = 0$$

$$r_y'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) r_y' + \frac{4q}{\gamma^3 m v_z^2} \left\langle y \frac{\partial \phi}{\partial y} \right\rangle \pm \frac{B'}{[B']} r_y - \frac{E_y^2}{v_y^3} = 0$$

(for electric focusing $\frac{B'}{[B']} \rightarrow \frac{q E'}{\gamma m v_z^2}$)

SPACE CHARGE TERM WITH ELLIPTICAL SYMMETRY

#4: ELLIPTICAL SYMMETRY: $\rho = \rho \left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2} \right)$

CAN BE SHOWN THAT $\langle x \frac{\partial \phi}{\partial x} \rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_x}{r_x + r_y}$

$\langle y \frac{\partial \phi}{\partial y} \rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_y}{r_x + r_y}$

DEFINING $Q = \frac{2q\lambda}{4\pi\epsilon_0 \gamma^3 m v_z^2}$

$$v_x'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) v_x' = \frac{2Q}{r_x + r_y} + \frac{B'}{[B_0]} v_x - \frac{\sigma_z^2}{r_x m} = 0$$

$$v_y'' + \frac{1}{\gamma v_z} \frac{d}{ds} (\gamma v_z) v_y' = \frac{2Q}{r_x + r_y} + \frac{B'}{[B_0]} v_y - \frac{\sigma_z^2}{r_y m} = 0$$

(for Electric Focusing $\frac{B'}{[B_0]} + \frac{qE'}{m\gamma v_z^2}$)



SPACE CHARGE TERN WITH ELLIPTICAL SYMMETRY II

J. BARNARD

(10)

ELLIPTICAL SYMMETRY:

$$\rho = \rho \left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2} \right)$$

CAN BE SHOWN THAT
(Sacherer, 1971)

$$\left\langle x \frac{\partial \Phi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_x}{r_x + r_y}$$

$$\left\langle y \frac{\partial \Phi}{\partial y} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_y}{r_x + r_y}$$

OUTLINE OF PROOF: (from R. Ryne)

$$\text{Let } \chi = \frac{x^2}{r_x^2 + s} + \frac{y^2}{r_y^2 + s}$$

$$\text{DEFINE } \eta(\chi) \text{ such that } \rho(x, y) = \frac{d\eta(\chi)}{d\chi} \Big|_{s=0} = \hat{\rho}(\chi) \Big|_{s=0}$$

$$\text{So } \rho = \hat{\rho} \left(\frac{x^2}{r_x^2} + \frac{y^2}{r_y^2} \right) = \hat{\rho}(\chi) \Big|_{s=0}$$

$$\text{DEFINE } \Phi(x, y) = \frac{-r_x r_y}{4\epsilon_0} \frac{\int_0^\infty \eta(\chi) ds}{\sqrt{r_x^2 + s} \sqrt{r_y^2 + s}}$$

$$\text{It follows that } \frac{\partial^2 \Phi}{\partial x^2} + \frac{\partial^2 \Phi}{\partial y^2} = -\frac{\rho}{\epsilon_0} \text{ AND SO IS A SOLUTION OF POISSON'S EQUATION (since } \Phi \rightarrow 0 \text{ as } x, y \rightarrow \infty)$$

WHAT IS $\left\langle x \frac{\partial \Phi}{\partial x} \right\rangle$?

$$\left\langle x \frac{\partial \Phi}{\partial x} \right\rangle = \frac{-r_x r_y}{4\pi\lambda\epsilon_0} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \rho(x, y) \int_0^\infty \frac{\eta' \frac{\partial \chi}{\partial x} ds}{\sqrt{r_x^2 + s} \sqrt{r_y^2 + s}}$$

$$\text{where } \lambda = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \rho(x, y)$$

So $\langle x \frac{\partial \phi}{\partial x} \rangle = \frac{-2V_x V_y}{4\lambda \epsilon_0} \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy x^2 \rho \left(\frac{x^2}{V_x^2} + \frac{y^2}{V_y^2} \right) \int_0^{\infty} \frac{\rho \left(\frac{x^2}{V_x^2 + s} + \frac{y^2}{V_y^2 + s} \right) ds}{(V_x^2 + s)^{3/2} (V_y^2 + s)^{3/2}}$

Let $r \cos \theta = \frac{x}{\sqrt{V_x^2 + s}}$ $r \sin \theta = \frac{y}{\sqrt{V_y^2 + s}}$

$\det J = \sqrt{V_x^2 + s} \sqrt{V_y^2 + s} r$ where J is the Jacobian
 $dx dy = \det J \cdot dr d\theta$

$\Rightarrow \langle x \frac{\partial \phi}{\partial x} \rangle = \frac{-2V_x V_y}{\lambda 2\epsilon_0} \int_0^{\infty} ds \int_0^{2\pi} d\theta \int_0^{\infty} dr r^3 \rho(r^2) \rho \left(\frac{V_x^2 + s}{V_x^2} r^2 \cos^2 \theta + \frac{V_y^2 + s}{V_y^2} r^2 \sin^2 \theta \right) \cos^2 \theta$

Let $r'^2 = \frac{V_x^2 + s}{V_x^2} r^2 \cos^2 \theta + \frac{V_y^2 + s}{V_y^2} r^2 \sin^2 \theta$

$= r^2 \left[1 + s \left(\frac{\cos^2 \theta}{V_x^2} + \frac{\sin^2 \theta}{V_y^2} \right) \right]$

with r fixed $2r' dr' = r^2 \left(\frac{\cos^2 \theta}{V_x^2} + \frac{\sin^2 \theta}{V_y^2} \right) ds$

$\Rightarrow \langle x \frac{\partial \phi}{\partial x} \rangle = \frac{-V_x V_y}{2\lambda \epsilon_0} \int_0^{\infty} ds \int_0^{2\pi} d\theta \int_r^{\infty} dr' \frac{2r' dr' r^3 \rho(r^2) \rho(r'^2) \cos^2 \theta}{r^2 \left(\frac{\cos^2 \theta}{V_x^2} + \frac{\sin^2 \theta}{V_y^2} \right)}$


$\int_0^{2\pi} \frac{\cos^2 \theta d\theta}{\frac{\cos^2 \theta}{V_x^2} + \frac{\sin^2 \theta}{V_y^2}} = \frac{2\pi V_x^2 V_y}{V_x + V_y}$

$\Rightarrow \langle x \frac{\partial \phi}{\partial x} \rangle = \frac{-V_x^3 V_y}{\lambda 2\pi \epsilon_0 (V_x + V_y)} \int_0^{\infty} dr 2\pi r \rho(r^2) \int_r^{\infty} dr' 2\pi r' \rho(r'^2)$

Recall: $\lambda = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} dx dy \rho(x, y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} dx dy \rho \left(\frac{x^2}{V_x^2} + \frac{y^2}{V_y^2} \right)$

Let $\frac{x}{V_x} = r \cos \theta$ $\frac{y}{V_y} = r \sin \theta$ $\det J = r V_x V_y$
 $\Rightarrow \lambda = \int_0^{\infty} \int_0^{2\pi} dr d\theta \rho(r^2) r V_x V_y = 2\pi r V_x V_y \int_0^{\infty} dr r \rho(r^2)$


$$\text{Now } \int_0^\infty dr r \hat{\rho}(r^2) \int_0^r dr' r' \hat{\rho}(r'^2) = \frac{1}{2} \int_0^\infty dr r \hat{\rho}(r^2) \int_0^\infty dr' r' \hat{\rho}(r'^2)$$



(By symmetry & consideration of diagram at left.)

$$\Rightarrow \left\langle x \frac{\partial \psi}{\partial x} \right\rangle = \frac{-\lambda}{4\pi\epsilon_0} \frac{r_x}{r_x + r_y}$$

CURRENT LIMIT FOR QUADRUPOLES



$$k = \begin{cases} \frac{B_0'}{CB(\beta)} & \text{MAGNETIC} \\ \frac{qE_0'}{\gamma m v_z^2} & \text{ELECTRIC} \end{cases}$$

$$r_x'' + k f(s) r_x - \frac{2Q}{r_x + r_y} = 0$$

$$r_y'' - k f(s) r_y - \frac{2Q}{r_x + r_y} = 0$$

(NOTE WE HAVE SET $\epsilon = 0$).

$$f(s) = \begin{cases} -1 & 0 < s < \pi L/2 \\ -1 & L - \pi L/2 < s < L + \pi L/2 \\ 1 & 2L - \pi L/2 < s < 2L \\ 0 & \text{otherwise} \end{cases}$$

$$\frac{1}{L} \int_0^{2L} f(s) \cos\left(\frac{\pi s}{L}\right) ds = \left(\frac{4}{\pi}\right) \sin\left(\frac{\pi \pi}{2}\right) = \text{fourier amplitude at fundamental lattice frequency}$$

$$\text{Let } r_x = r_b \left(1 + \delta \cos\left(\frac{\pi s}{L}\right)\right)$$

$$r_y = r_b \left(1 - \delta \cos\left(\frac{\pi s}{L}\right)\right)$$

$$\text{Let } k f(s) \rightarrow k \left(\frac{4}{\pi}\right) \sin\left(\frac{\pi \pi}{2}\right) \cos\left(\frac{\pi s}{L}\right)$$


COLLECTING "FAST" TERMS AND "SLOW" TERMS

$$\left[-\frac{\pi^2}{L^2} r_b \delta + k r_b \left(\frac{4 \sin(\frac{\pi \pi}{2})}{\pi}\right) \right] \cos\left(\frac{\pi s}{L}\right) = 0 \quad (\text{fast})$$

$$\delta k r_b \left(\frac{2 \sin(\frac{\pi \pi}{2})}{\pi}\right) = \frac{Q}{r_b} \quad (\text{slow})$$

$$\text{Fast } \Rightarrow \delta = \frac{4kL^2}{\pi^3} \sin\left(\frac{\pi \pi}{2}\right) \quad \& \quad Q_{\text{max}} \cong \frac{2\eta^2 k^2 L^2}{\pi^2} \left(\frac{\sin(\frac{\pi \pi}{2})}{(\frac{\pi \pi}{2})}\right)^2 r_b^2$$

CURRENT LIMIT FOR QUADRUPOLES



$$k = \begin{cases} \frac{B'_0}{CB(\beta)} & \text{MAGNETIC} \\ \frac{qE'_0}{\gamma m v_z^2} & \text{ELECTRIC} \end{cases}$$

$$r_x'' + k f(s) r_x - \frac{2Q}{r_x + r_y} = 0$$

$$r_y'' - k f(s) r_y - \frac{2Q}{r_x + r_y} = 0$$

(NOTE WE HAVE SET $\epsilon = 0$).

$$f(s) = \begin{cases} -1 & 0 < s < \pi L/2 \\ -1 & L - \pi L/2 < s < L + \pi L/2 \\ 1 & 2L - \pi L/2 < s < 2L \\ 0 & \text{otherwise} \end{cases}$$

$$\frac{1}{L} \int_0^{2L} f(s) \cos\left(\frac{\pi s}{L}\right) ds = \left(\frac{4}{\pi}\right) \sin\left(\frac{\pi \pi}{2}\right) = \text{fourier amplitude at fundamental lattice frequency}$$

$$\text{Let } r_x = r_b \left(1 + \delta \cos\left(\frac{\pi s}{L}\right)\right)$$

$$r_y = r_b \left(1 - \delta \cos\left(\frac{\pi s}{L}\right)\right)$$

$$\text{Let } k f(s) \rightarrow k \left(\frac{4}{\pi}\right) \sin\left(\frac{\pi \pi}{2}\right) \cos\left(\frac{\pi s}{L}\right)$$

COLLECTING "FAST" TERMS AND "SLOW" TERMS

$$\left[-\frac{\pi^2}{L^2} r_b \delta + k r_b \left(\frac{4 \sin(\frac{\pi \pi}{2})}{\pi}\right) \right] \cos\left(\frac{\pi s}{L}\right) = 0 \quad \text{(fast)}$$

$$\delta k r_b \left(\frac{2 \sin(\frac{\pi \pi}{2})}{\pi}\right) = \frac{Q}{r_b} \quad \text{(slow)}$$

$$\text{Fast } \Rightarrow \delta = \frac{4kL^2}{\pi^3} \sin\left(\frac{\pi \pi}{2}\right) \quad \& \quad Q_{\text{max}} \cong \frac{2\eta^2 k^2 L^2}{\pi^2} \left(\frac{\sin(\frac{\pi \pi}{2})}{(\frac{\pi \pi}{2})}\right)^2 r_b^2$$

Focusing term has both a fast and slow component:

$$\begin{aligned}kf(s)r_x &\rightarrow k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\cos\left(\frac{\pi s}{L}\right)r_b\left(1 + \delta\cos\left(\frac{\pi s}{L}\right)\right) \\ &= r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\cos\left(\frac{\pi s}{L}\right) + \delta r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\cos\left(\frac{\pi s}{L}\right)^2 \\ &= r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\cos\left(\frac{\pi s}{L}\right) + \delta r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\left(\frac{1}{2} + \frac{1}{2}\cos\left(\frac{2\pi s}{L}\right)\right) \\ &\cong r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\cos\left(\frac{\pi s}{L}\right) + \delta r_b k\left(\frac{4}{\pi}\right)\sin\left(\frac{\eta\pi}{2}\right)\left(\frac{1}{2}\right)\end{aligned}$$

CONTINUOUS FOCUSING

$$v_x'' = -k_{p0}^2 v_x + \frac{2Q}{v_x + v_y} - \frac{e^2}{v_x^2}$$

$$v_y'' = -k_{p0}^2 v_y + \frac{2Q}{v_x + v_y} - \frac{e^2}{v_y^2}$$

CURRENT LIMIT BALANCES PERVEANCE & EXTERNAL FOCUSING ($v_x = v_y = v_b$):

$$k_{p0}^2 v_b = \frac{Q_{max}}{v_b}$$

Effective k_{p0}^2 FOR QUADRUPOLES FOUND FROM DOMINANT FOURIER COMPONENT

$$k_{p0}^2 = \frac{2\eta^2 k^2 L^2}{\pi^2} \left(\frac{\sin(\frac{\eta\pi}{2})}{\frac{\eta\pi}{2}} \right)^2 \quad \text{where } k = \frac{B'}{B_0}$$

FOR CONTINUOUS FOCUSING: $k_{p0}^2 = \frac{\sigma_0^2}{4L^2}$

ELIMINATING L:

$$Q_{max} = \frac{\eta k \sigma_0}{\sqrt{2}\pi} \left(\frac{\sin \frac{\eta\pi}{2}}{\frac{\eta\pi}{2}} \right) v_b^2$$

← PERVEANCE LIMIT FOR FODO QUADRUPOLES

Envelope instabilities set upper limit on "single particle" phase advance σ_0




Experimental data (Tiefenback, 1986) from LBL's Single Beam Transport Experiment (SBTE)

79

Experimental limits on beam stability in terms of σ and σ_0

$\sigma_0 < 85^\circ$



SEE LUND & CHANLA 2006, NIMPR-A, FOR HIGHER ORDER PARTICLE-LATTICE RESONANCES WHICH CLAMPERS $\sigma_0 = 85^\circ$ LIMIT

SEE STRULLMENN & REISER, PARTICLE ACCELERATORS 14, 227, (1974) & LUND & BUCH, PLSTAB, I, 024801 (2004)

BACKWARD

QUADRUPOLE CURRENT LIMIT - CONTINUED

$$Q_{max} \approx \frac{\mu_0 \epsilon_0}{\sqrt{2\pi}} \left(\frac{\sin \frac{\pi}{2}}{\frac{\pi}{2}} \right) V_b^2$$

$$\text{here } k = \begin{cases} \frac{dB/dx}{[B]} \sim \frac{B}{[B] r_p} & \text{(MAGNETIC QUADROPOLE)} \\ \frac{q dE/dx}{\gamma m v_z^2} \sim \frac{z q V_q}{\gamma m v_z^2 r_p^2} & \text{(ELECTRIC QUADROPOLE)} \end{cases}$$

where $V_q = \frac{1}{2} \frac{dE}{dx} r_p^2$

So

$$Q_{max} \approx \frac{\mu_0 \epsilon_0}{\sqrt{2\pi}} \left(\frac{\sin \frac{\pi}{2}}{\frac{\pi}{2}} \right) \begin{cases} \frac{B}{[B]} \left[\frac{r_b}{r_p} \right] & \text{(MAGNETIC QUADROPOLE)} \\ \frac{z q V_q}{\gamma m v_z^2} \left[\frac{r_b^2}{r_p^2} \right] & \text{(ELECTRIC QUADROPOLE)} \end{cases}$$

Summary of Current Limits From Different Focusing Methods

EINZEL LENS

$$Q_{max} \approx \frac{3\pi^2}{8} \left(\frac{q b_0}{m v_0^2} \right)^2 \left(\frac{V_0}{L} \right)^2$$


SOLENOIDS

$$Q_{max} = \left(\frac{\omega_c V_0^2}{2\gamma \beta c} \right)^2$$

QUADROPOLE FOCUSING

$$Q_{max} \approx \frac{10^5}{\sqrt{2\pi}} \left(\frac{\sin \frac{\pi}{2}}{\frac{\pi}{2}} \right)$$

$$\left[\frac{B r_0}{C B P} \right] \left[\frac{2 q V_0}{\gamma m v_0^2} \right]$$



FOR NON-RELATIVISTIC BEAMS

$$\lambda_{max} \propto \frac{Q_0}{V}$$

$$\lambda_{max} \propto \frac{1}{m} B^2 r_p^2$$

$$\lambda_{max} \propto \left\{ \begin{array}{l} B_1 V_0^{1/2} r_p \\ V_0 \end{array} \right.$$

NOTE

Q_0 = Voltage between Einzel lenses

V_0 = Voltage on a grid relative to ground

V = particle energy / e