

Injector Simulations with warp

Daniel Winklehner

Simulation of Beam and Plasma Systems

D. Bruhwiler, R. Lehe, S. Lund, J.-L. Vay, D. Winklehner

USPAS, Old Dominion U., Hampton, VA, Jan 2018



Outline

- Morning:
 - Overview of ion sources
 - How extraction can be simulated in select cases
 - Sorted from “Easy” to “Hard” (very subjectively)
- Afternoon Lab I:
 - IBSimu crash course
 - Simulations of plasma ion sources using IBSimu
 - “Simple” plasma extraction + Adding B-field + Negative ions
- Afternoon Lab II:
 - Select challenges with low energy beam transport (LEBT)
 - Multiple species + space charge compensation
 - Warp simulations

Codes

- Codes (Raytracing/PIC + Plasma model):
 - IGUN (RZ) <http://www.egun-igun.com/>
 - IBSimu (RZ, 3D, 2D) <http://ibsimu.sourceforge.net/>
 - Warp (RZ, 3D, 2D) <http://warp.lbl.gov/>
 - Kobra-INP (RZ, 3D)
 -


Complications:

- Multiple ion species
- Added magnetic fields (see next):
 - Solenoid
 - Sextupole
- Negative ions/electrons
- 3D advisable!

Task 1

- Download https://people.nscl.msu.edu/~lund/uspas/sbp_2018/lec_inj/03.lebt_examples/solenoid_xy_slice.py
- Run it in warp
- Interpret the plot
- Add the second species (H₂⁺) by looking for “ASSIGNMENT Task 1” in the code and making the requested changes.
- What has changed? Did you change the current to account for two species?
- Now remove the protons and run again. Interpret!


Injection tests at BCS, Inc.








Versatile Ion Source (VIS) loaned from INFN

Cyclotron + Teststand provided by Best Cyclotron Systems, Inc. (BCS)

Simulations and Experiments Agree Well





Simulations and Experiments Agree Well



Space Charge Compensation

- Beam interacts with residual gas
- $$\sigma_e = \sigma_{ionization}$$
- $$\sigma_i = \sigma_{charge-exchange} + \sigma_{ionization}$$

$$\Delta\phi = \frac{I \cdot (1 - f_e)}{4\pi\epsilon_0 v_b}$$



Semi-Analytical Model

- 1975: Gabovich model for f_e , uses:
 - Secondary electron energy balance:

Steady state: energy transferred to electrons through Coulomb collisions = energy necessary to leave beam envelope

$$(\Delta\varphi_{neut})^2 = 3\mathcal{L} \cdot \frac{m_b}{m_e} \cdot \frac{\Phi_i}{U_0} \frac{n_b q e^2}{(4\pi\epsilon_0)^2} \left(\frac{q}{n_0\sigma_e} + \frac{v_b\sigma_i r_b}{2\bar{v}_i\sigma_e} \right)$$
$$f_e = 1 - \frac{\Delta\varphi_{neut}}{\Delta\varphi_{full}}$$

$$\Delta\varphi_{full} = \frac{I}{4\pi\epsilon_0 v_b} \quad \mathcal{L} = 4\pi \ln \left(4\pi\epsilon_0^{3/2} \frac{m_e^{3/2} v_b^3}{q e^3 n_e^{1/2}} \right)$$

M. Gabovich, L. Katsubo, and I. Soloshenko,
“Selfdecompensation of a stable quasineutral ion beam due to coulomb collisions”,
Fiz. Plazmy, vol. 1, pp. 304-309, 1975.

Semi-Analytical Model

- Major contributions to cross sections:

$$\sigma_e = \sigma_{ionization}$$

$$\sigma_i = \sigma_{charge-exchange} + \sigma_{ionization}$$

- Large uncertainties in available cross-section data!

- Other simplifications:

- Round, uniform beam
- Secondary ions: simple balance of produced ions = leaving ions
- Quasineutrality of the beam plasma

$$n_e = q \cdot n_b + n_i$$

$$(\Delta\varphi_{neut})^2 = 3\mathcal{L} \cdot \frac{m_b}{m_e} \cdot \frac{\Phi_i}{U_0} \frac{n_b q e^2}{(4\pi\epsilon_0)^2} \left(\frac{q}{n_0\sigma_e} + \frac{v_b\sigma_i r_b}{2\bar{v}_i\sigma_e} \right)$$

$$f_e = 1 - \frac{\Delta\varphi_{neut}}{\Delta\varphi_{full}}$$

Semi-Analytical Model


- Pressure in ECR transport line are as low as possible to reduce charge exchange (therefore low production of electrons)
- ECR beams are probably far from neutralized

$$n_e = q \cdot n_b + n_i \longrightarrow n_e = f_e \cdot (q \cdot n_b + n_i)$$




$$f_e = 1 - \sqrt{f_e} \cdot \frac{\Delta\varphi_{neut,Gabovich}}{\Delta\varphi_{full}}$$

$$\chi = \frac{\Delta\varphi_{neut,Gabovich}}{\Delta\varphi_{full}}$$


$$f_e = 1 + \frac{\chi^2}{2} - \frac{\chi}{2} \sqrt{\chi^2 + 4}$$




LEDA Injector Source



SuSI Beamline



f_e in SUSI beamline



Task 2

- Go back to using only protons in the python script. Now activate space charge compensation by looking for the line with “ASSIGNMENT Task 2” and uncommenting the `installbeforestep()` function.
- Run and interpret what you see. Save the figure
- After the run, calculate the mean f_e and use it instead of the function by commenting `installbeforestep` again and reducing the current appropriately.
- Run again and compare with the saved plot.