

your name _____

$$\begin{aligned}
 \vec{a} \times (\vec{b} \times \vec{c}) &= \vec{b}(\vec{a} \cdot \vec{c}) - \vec{c}(\vec{a} \cdot \vec{b}), \\
 \vec{a} \cdot (\vec{b} \times \vec{c}) &= \vec{b} \cdot (\vec{c} \times \vec{a}) = \vec{c} \cdot (\vec{a} \times \vec{b}), \\
 (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) &= (\vec{a} \cdot \vec{c})(\vec{b} \cdot \vec{d}) - (\vec{a} \cdot \vec{d})(\vec{b} \cdot \vec{c}), \\
 \nabla \times (\nabla \psi) &= 0, \\
 \nabla \cdot (\nabla \times \vec{a}) &= 0, \\
 \nabla \times (\nabla \times \vec{a}) &= \nabla(\nabla \cdot \vec{a}) - \nabla^2 \vec{a}, \\
 \nabla \cdot (\psi \vec{a}) &= \vec{a} \cdot \nabla \psi + \psi \nabla \cdot \vec{a}, \\
 \nabla \times (\psi \vec{a}) &= \nabla \psi \times \vec{a} + \psi \nabla \times \vec{a}, \\
 \nabla(\vec{a} \cdot \vec{b}) &= (\vec{a} \cdot \nabla) \vec{b} + (\vec{b} \cdot \nabla) \vec{a} + \vec{a} \times (\nabla \times \vec{b}) + \vec{b} \times (\nabla \times \vec{a}), \\
 \nabla \cdot (\vec{a} \times \vec{b}) &= \vec{b} \cdot (\nabla \times \vec{a}) - \vec{a} \cdot (\nabla \times \vec{b}), \\
 \nabla \times (\vec{a} \times \vec{b}) &= \vec{a}(\nabla \times \vec{b}) - \vec{b}(\nabla \times \vec{a}) + (\vec{b} \cdot \nabla) \vec{a} - (\vec{a} \cdot \nabla) \vec{b}, \\
 \nabla \cdot \vec{r} &= 3, \\
 \nabla \times \vec{r} &= 0, \\
 \nabla \cdot \hat{r} &= 2/r, \\
 \nabla \times \hat{r} &= 0, \\
 (\vec{a} \cdot \nabla) \hat{r} &= \frac{1}{r} [\vec{a} - \hat{r}(\vec{a} \cdot \hat{r})] = \frac{\vec{a}_\perp}{r}.
 \end{aligned}$$

$$\begin{aligned}
 \int_V d^3r \nabla \cdot \vec{A} &= \int_S d\vec{S} \cdot \vec{A}, \\
 \int_V d^3r \nabla \psi &= \int_S \psi d\vec{S}, \\
 \int_V d^3r \nabla \times \vec{A} &= \int_S d\vec{S} \times \vec{A}, \\
 \int_V d^3r (\phi \nabla^2 \psi + \nabla \phi \cdot \nabla \psi) &= \int_S \phi (d\vec{S}) \cdot \nabla \psi, \\
 \int_V d^3r (\phi \nabla^2 \psi - \psi \nabla^2 \phi) &= \int_S (\phi \nabla \psi - \psi \nabla \phi) \cdot d\vec{S}, \\
 \int_S (\nabla \times \vec{A}) \cdot d\vec{S} &= \oint d\vec{\ell} \cdot \vec{A}, \\
 \int_S (d\vec{S}) \times \nabla \psi &= \oint_C d\vec{\ell} \psi.
 \end{aligned}$$

$$\begin{aligned}
 \nabla^2 &= \partial_r^2 + \frac{2}{r} \partial_r - \frac{\ell(\ell+1)}{r^2}, \\
 \nabla^2 &= \partial_\rho^2 + \frac{1}{\rho} \partial_\rho - \frac{m^2}{r^2}, \\
 \nabla^2 \left(\frac{1}{r} \right) &= -4\pi \delta(\vec{r}).
 \end{aligned}$$

$$L^{\alpha}_{\beta} = \begin{bmatrix} \gamma & \gamma v & 0 & 0 \\ \gamma v & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

$$F_{\alpha\beta} = \begin{pmatrix} 0 & E_x & E_y & E_z \\ -E_x & 0 & -B_z & B_y \\ -E_y & B_z & 0 & -B_x \\ -E_z & -B_y & B_x & 0 \end{pmatrix},$$

$$F^{\alpha\beta} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix}.$$

$$m \frac{d}{dt} u^{\alpha} = e F^{\alpha\beta} u_{\beta},$$

$$\frac{d\vec{p}}{dt} = e\vec{E} + e\vec{v} \times \vec{B},$$

$$\omega_c = \frac{eB}{\gamma m}.$$

$$\nabla \cdot \vec{E} = 4\pi J^0,$$

$$(\nabla \times \vec{B}) - \partial_t \vec{E} = 4\pi \vec{J},$$

$$\nabla \cdot \vec{B} = 0,$$

$$\partial_t \vec{B} + \nabla \times \vec{E} = 0,$$

$$\partial_{\alpha} F^{\alpha\beta} = 4\pi J^{\beta},$$

$$\partial_{\alpha} \tilde{F}^{\alpha\beta} = 0.$$

$$e^2 = \frac{\hbar c}{137.036},$$

$$T^{\alpha\beta} = \pi^{\alpha} \partial^{\beta} \phi - g^{\alpha\beta} \mathcal{L},$$

$$\pi^{\alpha} \equiv \frac{\partial \mathcal{L}}{\partial(\partial_{\alpha} \phi)}$$

$$T^{00} = \frac{1}{8\pi} (E^2 + B^2),$$

$$T^{0i} = \frac{1}{4\pi} \epsilon_{ijk} E_j B_k,$$

$$T^{ij} = -T^i_j = \frac{1}{8\pi} (\delta_{ij} (E^2 + B^2) - 2E_i E_j - 2B_i B_j).$$

$$\vec{E} = -\nabla A_0 - \partial_t \vec{A} = -\nabla \Phi - \partial_t \vec{A}, \quad \vec{B} = \nabla \times \vec{A},$$

$$P_{\ell}(\cos \theta) = \sqrt{\frac{4\pi}{2\ell+1}} Y_{\ell m=0}(\theta),$$

$$Y_{0,0} = \frac{1}{\sqrt{4\pi}}, \quad Y_{1,0} = \sqrt{\frac{3}{4\pi}} \cos \theta,$$

$$Y_{1,\pm 1} = \mp \sqrt{\frac{3}{8\pi}} \sin \theta e^{\pm i\phi}, \quad Y_{2,0} = \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1),$$

$$Y_{2,\pm 1} = \mp \sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{\pm i\phi}, \quad Y_{2,\pm 2} = \sqrt{\frac{15}{32\pi}} \sin^2 \theta e^{\pm 2i\phi},$$

$$Y_{\ell-m}(\theta, \phi) = (-1)^m Y_{\ell m}^*(\theta, \phi),$$

$$\delta_{\ell\ell'} \delta_{mm'} = \int d\Omega Y_{\ell m}(\theta, \phi) Y_{\ell' m'}(\theta, \phi),$$

$$P_0(x) = 1, \quad P_1(x) = x,$$

$$P_2(x) = \frac{1}{2}(3x^2 - 1), \quad P_3(x) = \frac{1}{2}(5x^3 - 3x),$$

$$P_{\ell}(x=1) = 1, \quad \int_{-1}^1 dx P_{\ell}(x) P_{\ell'}(x) = \frac{2}{2\ell+1} \delta_{\ell\ell'},$$

$$(3D) \quad \Phi = \sum_{\ell m} (A_{\ell m} r^{\ell} + B_{\ell m} r^{-\ell-1}) Y_{\ell m}(\theta, \phi) e^{im\phi},$$

$$(2D) \quad \Phi = A_0 \ln(\rho) + \sum_m e^{im\phi} (A_m r^m + B_m r^{-m}),$$

$$\Phi = A_0 J_0 = \sum_m e^{im\phi} (A_m J_m(k\rho) + B_m N_m(k\rho)) e^{\pm kz},$$

$$\Phi = \frac{q}{r} + \frac{\vec{p} \cdot \vec{r}}{r^3} + \sum_m \frac{4\pi}{5r^3} q_{2m}(r) Y_{2m}(\theta, \phi),$$

$$\vec{E} = -\frac{1}{r^3} \vec{p} + 3 \frac{\vec{p} \cdot \vec{r}}{r^5} \vec{r} + \dots,$$

$$\Phi(r, \theta, \phi) = \sum_{\ell m} \frac{4\pi}{(2\ell+1)r^{\ell+1}} q_{\ell m}(r) Y_{\ell m}(\theta, \phi),$$

$$q_{22} = \sqrt{\frac{15}{32\pi}} \int d^3 r \rho(\vec{r}) (x - iy)^2 = \sqrt{\frac{15}{288\pi}} (Q_{11} - 2iQ_{12} - Q_{22}),$$

$$q_{21} = -\sqrt{\frac{15}{8\pi}} \int d^3 r \rho(\vec{r}) (x - iy)^2 = -\sqrt{\frac{15}{72\pi}} (Q_{13} - iQ_{23}),$$

$$q_{20} = \sqrt{\frac{5}{16\pi}} \int d^3 r \rho(\vec{r}) (3z^2 - r^2) = \sqrt{\frac{5}{16\pi}} Q_{33},$$

$$Q_{ij} \equiv \int d^3 r (3r_i r_j - r^2 \delta_{ij}) \rho(\vec{r}),$$

$$U = q\Phi_0 - \vec{p} \cdot \vec{E} - \frac{1}{6} Q_{ij} \partial_i \partial_j E_j,$$

$$A^\alpha(x) = \int d^4x' \frac{1}{|\vec{x} - \vec{x}'|} J^\alpha(x') \delta(x_0 - x'_0 - |\vec{x} - \vec{x}'|),$$

$$\vec{E} = e \left\{ \frac{\hat{n} \times [(\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}}]}{(1 - \vec{\beta} \cdot \hat{n})^3 |\vec{x}'|} \right\},$$

$$\vec{B} = \hat{n} \times \vec{E}.$$

$$P = \frac{2e^2}{3c} |\dot{\vec{\beta}}|^2 \text{ (Non-Rel.)},$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi(1 - \vec{\beta} \cdot \hat{n})^6} |(\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}}|^2,$$

$$P = \frac{2}{3c} e^2 \gamma^6 [\beta^2 - |\vec{\beta} \times \dot{\vec{\beta}}|^2],$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi(1 - \beta \cos\theta)^5} |\dot{\vec{\beta}}|^2 \sin^2\theta \text{ (linear)},$$

$$P = \frac{2e^2 \beta^2}{3c} \gamma^6 \text{ (linear)},$$

$$\frac{dP}{d\Omega} = \frac{e^2}{4\pi(1 - \beta n_\beta)^5} |\dot{\vec{\beta}}|^2 ((1 - \beta n_\beta)^2 - (1 - \beta^2) n_r^2) \text{ (circular)},$$

$$P = \frac{2}{3c} e^2 \beta^2 \gamma^4 \text{ (circular)}.$$

$$\frac{dP}{d\Omega} = \frac{1}{8\pi} k^2 |\hat{n} \times \vec{p}|^2,$$

$$P = \frac{\omega^4}{3} |\vec{p}|^2,$$

$$\text{(Thomson)} \quad \sigma = \frac{8\pi e^4}{3m^2},$$

$$\frac{\Delta\lambda}{\lambda} = \frac{\hbar\omega}{m} (1 - \cos\theta_s).$$

electron	-2.00231930436182 ± 0.0000000000000052
muon	-2.0023318418 ± 0.0000000013
proton	5.585694702 ± 0.000000017
neutron	-3.82608545 ± 0.00000090

$$\nabla^2 A^\alpha = -4\pi J^\alpha,$$

$$\vec{m} = \frac{1}{2} \int d^3r \vec{r} \times \vec{J} = \frac{I}{2} \int \vec{r} \times d\vec{l},$$

$$\vec{B} = -\frac{\vec{m}}{r^3} + \frac{3\vec{r}}{r^5} (\vec{m} \cdot \vec{r}),$$

$$\mu_e = \frac{e\hbar}{2m_e},$$

$$U = \frac{(\vec{\mu}_N \cdot \vec{\mu}_e)}{r^3} - \frac{3(\vec{\mu}_N \cdot \vec{r})(\vec{\mu}_e \cdot \vec{r})}{r^5} - \frac{8\pi}{3} (\vec{\mu}_N \cdot \vec{\mu}_e) \delta^3(\vec{r})$$

$$- \frac{e(\vec{\mu}_N \cdot \vec{L})}{mr^3},$$

$$T_{00} = \frac{1}{8\pi} (|\vec{E}|^2 + |\vec{B}|^2)$$

$$= \frac{a_i^2 + b_i^2}{8\pi} \frac{|\vec{a}|^2}{4\pi} \cos^2(\vec{k} \cdot \vec{r} - \omega t),$$

$$T_{0i} = \epsilon_{ijk} \frac{E_j B_k}{4\pi}$$

$$= \hat{k}_i \frac{|\vec{a}|^2}{4\pi} \cos^2(\vec{k} \cdot \vec{r} - \omega t),$$

$$T^{ij} = -T^i_j = \frac{1}{8\pi} (\delta_{ij}(E^2 + B^2) - 2E_i E_j - 2B_i B_j)$$

$$= \frac{1}{4\pi} \{ |\vec{a}|^2 \delta_{ij} - a_i a_j - b_i b_j \} \cos^2(\vec{k} \cdot \vec{r} - \omega t)$$

$$\omega_s = \omega \sqrt{\frac{1-v}{1+v}}$$

$$(TM) \quad E_z = \psi(x, y) e^{-i\omega t + ik_z z},$$

$$\Psi = 0 \text{ at boundary,}$$

$$\vec{E}_t(x, y) = \frac{ik_z}{(\omega^2 - k_z^2)} e^{-i\omega t + ik_z z} \nabla_t \psi(x, y),$$

$$\vec{B}_t(x, y) = \left(\frac{\omega}{k_z} \right) \hat{z} \times \vec{E}_t,$$

$$(TE) \quad B_z = \psi(x, y) e^{-i\omega t + ik_z z},$$

$$(\hat{n} \cdot \nabla_t) \psi(x, y)|_S = 0,$$

$$\vec{B}_t(x, y) = \frac{ik_z}{(\omega^2 - k_z^2)} e^{-i\omega t + ik_z z} \nabla_t \psi(x, y),$$

$$\vec{E}_t(x, y) = - \left(\frac{\omega}{k_z} \right) \hat{z} \times \vec{B}_t.$$

your name _____

LONG ANSWER SECTION

1. A thin wire extends from $x, y, z = (0, 0, -a)$ to $(0, 0, a)$. The current on the wire is wave-like

$$I = I_0 \cos(kz) \cos \omega_0 t, \quad k = \pi/2a.$$

- (a) (5 pts) Find the charge density on the wire as a function of time
- (b) (5 pts) What is the average power radiated by the wire
- (c) (5 pts) What is the angular distribution of the radiated power?

your name _____

Extra workspace for #1

your name _____

2. A square wave guide has transverse dimensions $0 < x < a$, $0 < y < a$. For a transverse magnetic wave (TM),
- (a) (10 pts) For a wave that propagates in the $+z$ direction with frequency ω , find a solution for the electric and magnetic fields, where the maximum electric field strength is E_0 . Choose the solution with the fewest transverse nodes.
 - (b) (5 pts) What is the lowest value of ω that leads to a propagating solution? And what is longitudinal velocity of the wave in that limit?

SHORT ANSWER SECTION

3. A linear accelerator accelerates either electrons or protons with an electric field of strength E_0 along the z direction. In terms of the ratio of masses M_p/M_e ,
- (a) (4 pts) Find the ratio of radiative powers, P_p/P_e , at the beginning of the acceleration when both particles would be moving non-relativistically.
 - (b) (4 pts) Find the ratio of radiative powers, P_p/P_e , at the end of the acceleration when both particles would be moving ultra-relativistically with energy K .
4. (4 pts) Two gases of fully ionized particles are stored in identical containers with identical numbers of ions (and electrons). In gas A , the gas is fully ionized hydrogen, where all the positive ions are protons. In gas B , the positive ions are deuterons (one neutron and one proton), which have approximately twice the mass of a proton. The number of free electrons in each container is thus equal. The mean free paths, ℓ_A and ℓ_B , of light traveling through the containers vary by approximately what factor? Circle the correct answer:

$$\ell_A/\ell_B \approx$$

- (a) $\frac{1}{16}$
- (b) $\frac{1}{8}$
- (c) $\frac{1}{4}$
- (d) $\frac{1}{2}$
- (e) 1
- (f) 2
- (g) 4
- (h) 8
- (i) 16