


# Nuclear properties that are relevant for reaction rates:

Nucleons in the nucleus can only have discrete energies. Therefore, the nucleus as a whole can be excited into discrete energy levels (excited states)



Each state is characterized by:

- energy (mass)
- spin
- parity
- lifetimes against  $\gamma$ , p, n, and  $\alpha$  emission

The **lifetime** is usually given as a width as it corresponds to a width in the excitation energy of the state according to Heisenberg:

$$\Delta E \cdot \Delta t = \hbar$$


therefore, a lifetime  $\tau$  corresponds to a width  $\Gamma$ :

$$\Gamma = \frac{\hbar}{\tau}$$

the lifetime against the individual “channels” for  $\gamma$ , p, n, and  $\alpha$  emission are usually given as **partial widths**

$$\Gamma_{\gamma}, \Gamma_p, \Gamma_n, \text{ and } \Gamma_{\alpha} \quad \text{with} \quad \Gamma = \sum \Gamma_i$$


# A Real Example



# Basic reaction mechanisms involving strong or electromagnetic interaction:

Example: neutron capture  $A + n \rightarrow B + \gamma$


## I. Direct reactions (for example, direct capture)




direct transition into bound states

## II. Resonant reactions (for example, resonant capture)

Step 1: Compound nucleus formation  
(in an unbound state)




Step 2: Compound nucleus decay




or a resonant  $A(n,\alpha)B$  reaction:

Step 1: Compound nucleus formation  
(in an unbound state)



Step 2: Compound nucleus decay




For resonant reactions,  $E_n$  has to “match” an excited state (but all excited states have a width and there is always some cross section through tails)

But enhanced cross section for  $E_n \sim E_x - S_n$

more later ...

# Direct reactions - for example direct capture:



Direct transition from initial state  $|a+A\rangle$  to final state  $\langle f|$  (some state in B)

$$\sigma \propto \pi \lambda_a^2 \cdot \left| \langle f | H | a + A \rangle \right|^2 \cdot P_l(E)$$

geometrical factor  
(deBroglie wave length  
of projectile - "size" of  
projectile)

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2mE}}$$

Interaction matrix  
element

Penetrability: probability  
for projectile to reach  
the target nucleus for  
interaction.


Depends on projectile  
Angular momentum  $l$   
and Energy  $E$

$$\sigma \propto \frac{1}{E} \cdot \left| \langle f | H | a + A \rangle \right|^2 \cdot P_l(E)$$

III.25

Penetrability: 2 effects that can strongly reduce penetrability:

1. Coulomb barrier



for a projectile with  $Z_2$  and a nucleus with  $Z_1$

$$V_c = \frac{Z_1 Z_2 e^2}{R} \quad \text{or} \quad V_c [\text{MeV}] = 1.44 \frac{Z_1 Z_2}{R[\text{fm}]} \approx 1.2 \frac{Z_1 Z_2}{(A_1^{1/3} + A_2^{1/3})}$$

Example:  $^{12}\text{C}(p,\gamma)$   $V_c = 3 \text{ MeV}$


Typical particle energies in astrophysics are  $kT = 1\text{-}100 \text{ keV}$  !

Therefore, all charged particle reaction rates in nuclear astrophysics occur way below the Coulomb barrier – fusion is only possible through tunneling

## 2. Angular momentum barrier

Incident particles can have orbital angular momentum  $L$

Classical:



Momentum  $p$

Impact parameter  $d$

$$L = pd$$

In quantum mechanics the angular momentum of an incident particle can have discrete values:

$$L = \sqrt{l(l+1)} \hbar$$

With

$$l = 0$$

s-wave

And parity of the

$$l = 1$$

p-wave

wave function:  $(-1)^l$

$$l = 2$$

d-wave

...

For radial motion (with respect to the center of the nucleus), angular momentum conservation (central potential !) leads to an energy barrier for non zero angular momentum.

Classically, one needs the radial kinetic energy to overcome the central potential, but if  $d \neq 0$  then there is an increasing amount of “non radial kinetic energy”, which one needs to supply as well (at  $z=0$  for example,  $K_r=0$ , but of course  $K \neq 0$ ) 8



Energy  $E$  of a particle with angular momentum  $L$  (still classical)

$$E = \frac{L^2}{2mr^2}$$

Similar here in quantum mechanics:

$$V_l = \frac{l(l+1)\hbar^2}{2\mu r^2}$$

$\mu$  : reduced mass of projectile-target system

Peaks again at nuclear radius (like Coulomb barrier)

Or in MeV using the nuclear radius and mass numbers of projectile  $A_1$  and target  $A_2$ :

$$V_l [\text{MeV}] = 12 \frac{l(l+1)}{\left( \frac{A_1 A_2}{A_1 + A_2} \right) (A_1^{1/3} + A_2^{1/3})}$$


## Direct reactions – the simplest case: s-wave neutron capture

No Coulomb or angular momentum barriers:  $V_l=0$

$$V_C=0$$

s-wave capture therefore always dominates at low energies

But, change in potential still causes reflection – even without a barrier  
Recall basic quantum mechanics:



Transmission proportional to  $\sqrt{E}$


Therefore, for direct s-wave neutron capture:

Penetrability  $P_l(E) \propto \sqrt{E}$


Cross section (use Eq. III.19):  $\sigma \propto \frac{1}{\sqrt{E}}$  Or  $\sigma \propto \frac{1}{v}$

### Example: ${}^7\text{Li}(n,\gamma)$

thermal  
cross section  
 $\langle\sigma\rangle=45.4\text{ mb}$   
(see Pg. 27)



## Why s-wave dominated ? Level scheme:



## Angular momentum and parity conservation:

Entrance channel  ${}^7\text{Li} + n$  :  $3/2^- + 1/2^+ + l^{(-1)^l} = 1^-, 2^-$  ( $l=0$  for s-wave)

Exit channel  ${}^8\text{Li} + \gamma$  :  $2^+ + ?$  (photon spin/parity)

Recall: Photon angular momentum/parity depend on multipolarity:

For angular momentum  $L$  (=multipolarity) electric transition  $E_L$  parity  $(-1)^L$   
magnetic transition  $M_L$  parity  $(-1)^{L+1}$

Also recall:

E.M. Transition strength increases:

- for lower L
- for E over M
- for higher energy  $\propto E_\gamma^{2L+1}$

Entrance channel  ${}^7\text{Li} + n$  :  $3/2^- + 1/2^+ + l^{(-)}$  =  $1^-, 2^-$  (  $l=0$  for s-wave )

Exit channel  ${}^8\text{Li} + \gamma$  :  $2^+ + 1^-$  =  $1^-, 2^-, 3^-$

E1 photon  
lowest EL that  
allows to fulfill  
conservation laws

↑  
match possible

Same for  $1^+$  state

→ “At low energies  ${}^7\text{Li}(n,\gamma)$  is dominated by (direct) s-wave E1 capture”.

## Stellar reaction rate for s-wave neutron capture:

Because  $\sigma \propto \frac{1}{v} \longrightarrow \sigma v = \text{const} = \langle \sigma v \rangle$

## Direct reactions – neutron captures with higher orbital angular momentum

For neutron capture, the only barrier is the angular momentum barrier

The penetrability scales with

$$P_l(E) \propto E^{1/2+l}$$

and therefore the cross section (Eq III.19)


$$\sigma \propto E^{l-1/2}$$

for  $l > 0$  cross section decreases with decreasing energy (as there is a barrier present)

Therefore, s-wave capture in general dominates at low energies, in particular at thermal energies. Higher  $l$ -capture usually plays only a role at higher energies. What “higher” energies means depends on case to case - sometimes s-wave is strongly suppressed because of angular momentum selection rules (as it would then require higher gamma-ray multipolarities)

Example: p-wave capture in  $^{14}\text{C}(n,\gamma)^{15}\text{C}$


$$\sigma \propto \sqrt{E}$$



(from Wiescher et al. ApJ 363 (1990) 340)



## Why p-wave ?



### Exit channel ( $^{15}\text{C} + \gamma$ )

	$\gamma$	total to $1/2^+$	total to $5/2^+$	
<b>E1</b>	<b><math>1^-</math></b>	<b><math>1/2^- \quad 3/2^-</math></b>	<b><math>3/2^- \quad 5/2^- \quad 7/2^-</math></b>	<b>strongest !</b>
M1	$1^+$	$1/2^+ \quad 3/2^+$	$3/2^+ \quad 5/2^+ \quad 7/2^+$	
E2	$2^+$	$3/2^+ \quad 5/2^+$	$1/2^+ \quad 3/2^+ \quad 5/2^+ \quad 7/2^+ \quad 9/2^+$	

### Entrance channel:

	$ \pi$	$^{14}\text{C}$	$n$	total	strongest possible Exit multipole	
					into $1/2^+$	into $5/2^+$
s-wave	$0^+$	$0^+$	$1/2^+$	$1/2^+$	M1	E2
p-wave	$1^-$	$0^+$	$1/2^+$	$1/2^- \quad 3/2^-$	<b>E1</b>	<b>E1</b>

despite of higher barrier, for relevant energies (1-100 keV) p-wave E1 dominates. At low energies, for example thermal neutrons, s-wave still dominates. But here for example, the thermal cross section is exceptionally low ( $<1\mu\text{b}$  limit known)

# Charged particle induced direct reactions

## Cross section and S-factor definition

(for example proton capture - such as  $^{12}\text{C}(p,\gamma)$  in CN cycle)

incoming projectile  $Z_1 A_1$  (for example proton or  $\alpha$  particle)  
target nucleus  $Z_2 A_2$

again

$$\sigma \propto \frac{1}{E} \cdot P_l(E) \cdot \left| \langle f | H | a + A \rangle \right|^2$$


but now incoming particle has to overcome Coulomb barrier. Therefore

$$P_l(E) \propto e^{-2\pi\eta} \quad \text{with} \quad \eta = \sqrt{\frac{\mu}{2E}} \frac{Z_1 Z_2 e^2}{\hbar}$$


(from basic quantum mechanical barrier transmission coefficient)

# Penetrability factor $P_l(E)$ example

Charged particle (proton)



Neutron



(from Iliadis “Nuclear Physics of Stars”)

## The concept of the astrophysical S-factor (for n-capture)

recall:

$$\sigma \propto \underbrace{\frac{1}{E} \cdot P_l(E)}_{\text{“trivial” strong energy dependence}} \cdot \underbrace{\left| \langle f | H | a + A \rangle \right|^2}_{\text{“real” nuclear physics weak energy dependence (for direct reactions !)}} \quad \text{III.25}$$

S-factor concept: write cross section as

**strong “trivial” energy dependence      X      weakly energy dependent S-factor**

The S-factor can be

- easier graphed
- easier fitted and tabulated
- easier extrapolated
- and contains all the essential nuclear physics

Note: There is no “universally defined S-factor - the S-factor definition depends on the type of reaction and (for neutrons at least) on l-value

Here the main energy dependence of the cross section (for direct reactions !)  
is given by

$$\sigma \propto \frac{1}{E} e^{-\frac{b}{\sqrt{E}}}$$

$$b = 31.28 \cdot Z_1 Z_2 A^{1/2} \sqrt{\text{keV}}$$

$$A = \frac{A_1 A_2}{A_1 + A_2} = \frac{\mu}{m_U}$$

therefore the **S-factor for charged particle reactions is defined** via


$$\sigma = \frac{1}{E} e^{-b/\sqrt{E}} S(E)$$

typical unit for  $S(E)$ : keV barn

So far this all assumed s-wave capture. However, the additional angular momentum barrier leads only to a roughly constant addition to this S-factor that strongly decreases with  $l$

Therefore, the S-factor for charged particle reactions is defined independently of the orbital angular momentum


Example:  
 $^{12}\text{C}(p,\gamma)$  cross section



**Fig. 4-4** The measured cross section for the reaction  $\text{C}^{12}(p,\gamma)\text{N}^{13}$  as a function of laboratory proton energy. A four-parameter theoretical curve has been fitted to the experimental points. An extrapolation to  $E_p = 0.025$  Mev, which is an interesting energy for this reaction in astrophysics, appears treacherous. (Courtesy of W. A. Fowler and J. L. Vogl.)

## S-Factor:

Need rate  
about here




From the **NACRE compilation** of charged particle induced reaction rates on stable nuclei from H to Si (Angulo et al. Nucl. Phys. A 656 (1999) 3)

## 9.3.2. Relevant cross section - Gamov Window

for charged particle reactions

$$\langle \sigma v \rangle = \sqrt{\frac{8}{\pi\mu}} (kT)^{-3/2} \int \sigma(E) E e^{-\frac{E}{kT}} dE = \sqrt{\frac{8}{\pi\mu}} (kT)^{-3/2} \int S(E) e^{-\left(\frac{b}{\sqrt{E}} + \frac{E}{kT}\right)} dE$$

**Gamov Peak**



Note: relevant cross section in tail of M.B. distribution, much larger than  $kT$  (very different from n-capture !)



The Gamov peak can be approximated with a Gaussian

$$e^{-\frac{b}{\sqrt{E}} - \frac{E}{kT}} \quad e^{-\frac{3E_0}{kT}} \quad e^{-\frac{E - E_0}{E/2}}^2$$

centered at same energy  $E_0$  with width  $\Delta E$  such that  $d^2/dE^2|_{E_0}$  is the same


Then, the **Gamov window** or the range of relevant cross section can be easily calculated using:

$$E_0 = \left( \frac{bkT}{2} \right)^{3/2} = 0.12204 (Z_1^2 Z_2^2 A)^{1/3} T_9^{2/3} \text{ MeV}$$

$$\Delta E = \frac{4}{\sqrt{3}} \sqrt{E_0 kT} = 0.23682 (Z_1^2 Z_2^2 A)^{1/6} T_9^{5/6} \text{ MeV}$$

with A “reduced mass number” and  $T_9$  the temperature in GK

Example:



**Fig. 4-7** The Gamow peak for the reaction  $C^{12}(p, \gamma)N^{13}$  at  $T = 30 \times 10^6 \text{ K}$ . The curve is actually somewhat asymmetric about  $E_0$ , but it is nonetheless adequately approximated by a gaussian.

### 9.3.3. Reaction rate from S-factor

Often (for example with theoretical reaction rates) one approximates the rate calculation by assuming the S-factor is constant over the Gamov Window

$$S(E)=S(E_0)$$

then one finds the useful equation:

$$N_A \langle \sigma v \rangle = 7.83 \cdot 10^9 \left( \frac{Z_1 Z_2}{A T_9^2} \right)^{1/3} S(E_0) [\text{MeV barn}] e^{-4.2487 \left( \frac{Z_1^2 Z_2^2 A}{T_9} \right)^{1/3}}$$

Equation III.53

(A reduced mass number !)

better (and this is often done for experimental data) one expands  $S(E)$  around  $E=0$  as powers of  $E$  to second order:

$$S(E) = S(0) + ES'(0) + \frac{1}{2} ES''(0)$$

If one integrates this over the Gamov window, one finds that one can use Equation III.46 when **replacing  $S(E_0)$  with the effective S-factor  $S_{eff}$**

$$S_{eff} = S(0) \left[ 1 + \frac{5}{12\tau} + \frac{S'(0)}{S(0)} \left( E_0 + \frac{35}{36} kT \right) + \frac{1}{2} \frac{S''(0)}{S(0)} \left( E_0^2 + \frac{89}{36} E_0 kT \right) \right]$$

with  $\tau = \frac{3E_0}{kT}$  and  $E_0$  as location of the Gamov Window (see Pg. 51)