

Global properties of atomic nuclei


How to probe nuclear size?

⇒ Electron Scattering from nuclei

For low energies and under conditions where the electron does not penetrate the nucleus, the electron scattering can be described by the Rutherford formula. The *Rutherford formula* is an analytic expression for the differential scattering cross section, and for a projectile charge of e , it is


$$\left(\frac{d\sigma}{d\cos\theta} \right)_R = \frac{\pi}{2} \left[\frac{\hbar c Z \alpha}{T_{KE} (1 - \cos\theta)} \right]^2$$

← Kinetic energy of electron



As the energy of the electrons is raised enough to make them an effective nuclear probe, a number of other effects become significant, and the scattering behavior diverges from the Rutherford formula. The probing electrons are relativistic, they produce significant nuclear recoil, and they interact via their magnetic moment as well as by their charge. When the magnetic moment and recoil are taken into account, the expression is called the *Mott cross section*.

A major period of investigation of nuclear size and structure occurred in the 1950's with the work of Robert Hofstadter and others who compared their high energy electron scattering results with the Mott cross section. The illustration below from Hofstadter's work shows the divergence from the Mott cross section which indicates that the electrons are penetrating the nucleus - departure from point-particle scattering is evidence of the structure of the nucleus.



The cross section from elastic electron scattering is:

$$\frac{d\sigma}{d \cos \theta} = \left(\frac{d\sigma}{d \cos \theta} \right)_{\text{point}} |F(q)|^2$$


Mott cross section **form factor**

Form factor

$$F(\vec{q}) = \int d^3r \rho_{\text{ch}}(\vec{r}) e^{i\vec{q}\vec{r}}$$

q – three momentum transfer of electron

Sizes




$$\rho(0) = 0.16 \text{ nucleons/fm}^3$$

$$\rho(r) = \rho_0 \left[1 + \exp\left(\frac{r - R}{a}\right) \right]^{-1}$$


$$R \approx 1.2A^{1/3} \text{ fm}, \quad a \approx 0.6 \text{ fm}$$

Calculated and measured densities




Protons and neutrons aren't point particles

charge distribution in the neutron



charge distribution in the proton



mass →	2.4 MeV	1.27 GeV	171.2 GeV
charge →	2/3	2/3	2/3
spin →	1/2	1/2	1/2
name →	u up	c charm	t top
Quarks	4.8 MeV	104 MeV	4.2 GeV
	-1/3	-1/3	-1/3
	1/2	1/2	1/2
	d down	s strange	b bottom

$$\langle R_{\text{ch}}^2 \rangle = \langle R_{\text{pp}}^2 \rangle + \langle r_{\text{p}}^2 \rangle + \frac{N}{Z} \langle r_{\text{n}}^2 \rangle + \frac{3\hbar^2}{4m_{\text{p}}^2 c^2}$$

$$\sqrt{\langle r_{\text{p}}^2 \rangle} = 0.8775(51) \text{ fm}$$

$$\langle r_{\text{n}}^2 \rangle = -0.1149(27) \text{ fm}^2$$

relativistic Darwin-Foldy correction

Proton size puzzle

<http://www.newscientist.com/article/dn19141-incredible-shrinking-proton-raises-eyebrows.html>


<https://www.psi.ch/media/proton-size-puzzle-reinforced>

Muon has a mass of 105.7 MeV, which is about 200 times that of the electron


Bohr radius:
$$a_0 = \frac{\hbar}{m_e c \alpha}$$

New radius: 0.84087(39) fm

Proton radius determinations over time





Proton charge radii obtained from hydrogen spectroscopy



New Measurement Deepens Proton Puzzle

Aug. 2016: Pohl *et al.* (Science 353) determined the charge radius of the deuteron, a nucleus consisting of a proton and a neutron, from the transition frequencies in muonic deuterium. Mirroring the proton radius puzzle, the radius of the deuteron was several standard deviations smaller than the value inferred from previous spectroscopic measurements of electronic deuterium. This independent discrepancy points to experimental or theoretical error or even to physics beyond the standard model.



Examples of how the measured values of constants can vary dramatically before converging on their correct values (from PDG)